

# Abstracts of the International Congress of Future Medical Pioneers 2021

ICOFMEP 2021  
8-9 May 2021, Virtual Event  
www.futuremedicalpioneers.org

## Scientific Programme Committee Chair(s):

Professor Bahar Uslu, PhD	Quinnipiac University Frank H. Netter School of Medicine	Hamden, CT, US
Assoc. Prof. Ahmet Şen	Health Sciences University, Samsun Training and Research Hospital	Samsun, Turkey

## Organizational Board

Professor Bahar Uslu, PhD	Quinnipiac University Frank H. Netter School of Medicine	Hamden, CT, US
Assoc. Prof. Ahmet Şen	Health Sciences University, Samsun Training and Research Hospital	Samsun, Turkey
Assoc. Prof. Enes Emre Başar	Anadolu University, Faculty of Business Administration	Eskişehir, Turkey
Kenan Demir, MD	Health Sciences University, Samsun Training and Research Hospital	Samsun, Turkey
Özlem Sezer, MD	Health Sciences University, Samsun Training and Research Hospital	Samsun, Turkey

## Abstract Review Panel

Each abstract was scored double blind review.

Fevzi Toraman	Fatih Kılıçbay	Gökşen Derya Reis Köse
Handan Çelik	Murat Serkant Ünal	Gülsemir Çiçek
İmer Okar	Sude Hatun Aktimur	Kenan Demir
Mehmet Bilge Çetinkaya	Aykut Öztürk	Mehmet Cenk Turgut
Ünal Uslu	Bahadır Yazıcıoğlu	Onur Taşcı
Başak Büyük	Betül Sarı	Özlem Sezer
Başar Erdivanlı	Büşra Akpınar	Saim Gubari
Deniz Aka Satar	Canan Soyer Çalışkan	Samettin Çelik
Mahcubey Çubukçu	Eda Türe	Selim Görgün
Özgür Günel	Elif Hilal Ünverdi	Şadiye Açıkgöz
Pınar Tosun Taşar	Emine Aksoy	Yasemin Nasır
Aysın Türkmen	Emine Tural	Murat Kubat
Emine Özçınar	Göksenin Ünlügüzel Üstün	

## Table of Contents

Lockdown effect on emergency surgical consults during the covid-19 outbreak: Experience of a pandemic hospital	
Ahmet Burak Ciftçi and Sönmez Ocak .....	<b>1</b>
Fusion Planet: Past, Today and Future of Computer Scientific Solutions in Digital Pathology	
Taha Yiğit Alkan, Hüseyin Gökhan Akçay and Havva Serap Toru.....	<b>2</b>
14q32.31q32.33 Microdeletion Detected by Chromosomal Microarray in a Child with Dysmorphism and Hypotonia	
Aslıhan Sanrı and Özlem Sezer.....	<b>4</b>
Factors Associated with Pressure Ulcers in Adult Critical Care Patients	
Başar Erdivanlı.....	<b>5</b>
COVID-19 Through the Eyes of Physiatrists: Back Pain Can be a Symptom and May Predict Pneumonia in COVID-19	
Cuma Uz.....	<b>6</b>
The Effect of Ulipristal Acetate and Vitamin D3 on Folliculogenesis	
Damla Gül Fındık, Gülnur Take Kaplanoğlu, Gökçe Nur Arık and Nagva B. Abubaker Alemari ....	<b>7</b>
Evaluation of Antibody Levels in Pcr Negative Covid-19 Suspected Case Series	
Demet Gür Vural,Büşra Usta, Yeliz Tanrıverdi Çaycı, Kemal Bilgin and Asuman Birinci.....	<b>8</b>
Evaluation of Antinuclear Antibodies in Patients who were Infected with Covid 19	
Demet Gür Vural .....	<b>9</b>
Depression, Hopelessness, and Social Support in Hemodialysis Atients	
Demet Yavuz.....	<b>11</b>
Effects of thymoquinone on anxiety-related behavior and auditory potentials in rats exposed to glucocorticoid	
Deniz Kantar .....	<b>12</b>
Spontaneous Tumor Lysis Syndrome in Lung Cancer: Very Rare Case for Solid Tumors	
Buğra Özel, Düriye Sıla Karagöz Özen and Ahmet Baran .....	<b>13</b>
Is Selenium deficiency related to an increased risk for diabetic retinopathy?	
Hacer Pınar Öztürk Kurt, Düriye Sıla Karagöz Özen, İpek Genç, Mukadder Erdem and Mehmet Derya Demirağ .....	<b>14</b>
The Development of Family Medicine Identity Scale	
Duygu Üstünoğlu, İsmail Kasım and Adem Özkara.....	<b>15</b>
Investigation of the relationship between magnesium level and vitamin D, bone mineral density, knee osteoarthritis and chronic diseases	
Ebru Yılmaz and Sena Ünver.....	<b>16</b>
Outcomes of Breast Cancer Brain Metastasis Patients Undergoing Stereotactic Radiotherapy	
Ela Delikgöz Soykut and Nilgün Şahin.....	<b>17</b>
Epithelial-mesenchymal transition and Cancer	
Elif Önder .....	<b>19</b>
Ethical Aspects of Embryo Research	
E. Elif Vatanoğlu-Lutz .....	<b>20</b>
From Tuhfetü'l Tıb* to the Medicine of the Future	
Eser Epözdemir, E. Elif Vatanoğlu-Lutz and Bahar Uslu.....	<b>21</b>

Medically Assisted Reproduction in Turkish Law	
Emel Badur .....	<b>22</b>
The Cytotoxicity Effects of Metformin and Lithium Substances in the Human Intestinal Caco-2 Cells	
Emine Tural.....	<b>23</b>
A Comparison of Propofol with Ketofol For Sedation Quality and Side Effects in Patients Undergoing Colonoscopy	
Ender Çam, Deniz Karakaya and Sibel Barış .....	<b>24</b>
International scientific collaborative activities: Barriers and opportunities	
Şule Başar and Enes Emre Başar.....	<b>25</b>
Clinical outcomes of intraarticular PRP and corticosteroid combination in advanced osteoarthritis	
Erhan Okay .....	<b>27</b>
A COVID-19 case complicated by ecchymosis and cyanosis	
Esmeray Mutlu Yılmaz .....	<b>28</b>
The frequency of micronodular type neuroendocrine cell hyperplasia detected in gastric endoscopic biopsies and its relationship with clinicopathological parameters	
Oğuzhan Okcu, Ezgi Hacıhasanoğlu and Bayram Şen .....	<b>29</b>
Renal Tissue Morphology and Morphometric Alterations with Different Fixatives (Formalin, Bouin's, Alfac and B5 Solutions)	
Fatma Mert, Bilge Serdaroğlu, İbrahim Alptekin, Ferda Topal Çelikkan and Oya Evirgen .....	<b>30</b>
New challenge in adolescents: 'at risk polycystic ovary syndrome' estimation	
Fatma Nurgül Taşgöz.....	<b>31</b>
Determination of Cell Viability in LNCaP Prostate Cancer Cells Treated with Boric Acid	
Gizem Kabasakal and Mücahit Seçme.....	<b>33</b>
Comparison of Insulin Treatment with Mesenchymal Stem Cell Treatment in Experimental Type 1 Diabetes-Induced Rats	
Gökçen Gökçe, Murat Tosun, Hasan Hüseyin Demirel and Esra Aslan .....	<b>34</b>
Oxidative Effects and Biochemical Markers Related to Disease Severity Parameters in Cystic Fibrosis Cases	
Gökseven Ünlügüzel Üstün.....	<b>35</b>
Determination of Sleep Quality in Patients with Rotator Cuff Tears	
Gonca Sağlam .....	<b>37</b>
Determination of the functions of miRNAs associated with PI3K-Akt signaling pathway and p53 signaling pathway in prostate cancer	
Gözde Öztan.....	<b>38</b>
Efficacy of Lacosamide Therapy in Focal Onset Refractory Epilepsy of Childhood: A Single Center Experience	
Halil Ural Aksoy, Celil Yılmaz, Senem Ayca and Muzaffer Polat.....	<b>39</b>
New treatment options in rare diseases	
İşıl Özer .....	<b>40</b>
Evaluation of the Effects of CoenzymeQ10 Treatment on Corneal Epithelium by VEGF and VEGFR in STZ-induced Diabetes Rats	
Çiğdem Karaca, Müberra Akdoğan and Ruhi Türkmen.....	<b>41</b>
Investigation of Colistin Resistance in Gram Negative Bacteria by Colistin Susceptibility Tube Test	
Kübra Hacıeminoğlu Ülker, Yeliz Tanriverdi Çaycı, Eda Köprü and Asuman Birinci .....	<b>42</b>

Early Detection of Antibiotic Susceptibility of Enterobacteriaceae Family Isolates Using an Inverted Microscope; Preliminary Study	
Kemal Bilgin, Yeliz Tanrıverdi Çaycı, Demet Gür Vural and Asuman Birinci.....	<b>43</b>
Optimization Study of Detecting Antibiotic Sensitivity of Pseudomonas aeruginosa Using an Inverted Microscope	
Kemal Bilgin.....	<b>45</b>
The distribution of Enterobacteriales isolated from the urine samples of children, and the evaluation of antimicrobial susceptibility	
Yeliz Tanrıverdi Çaycı, Gülşah Karacan, Moein Yoosefi, Kemal Bilgin, Demet Gür Vural and Asuman Birinci.....	<b>47</b>
Future Expectations in ART	
İlay Gözükara and M. Turan Çetin.....	<b>49</b>
Can We Consider Embryos Solely as Biological Material?	
Maide Barış.....	<b>50</b>
Common Chest CT Findings in 100 COVID-19 Patients Followed for Pneumonia	
Mehmet Akçiçek.....	<b>51</b>
Does COVID-19 infection trigger the formation of Anti-nuclear Antibodies?	
Melek Bilgin and Eşe başbulut.....	<b>52</b>
Guillain Barre Syndrome Induced by Covid 19: Discussion with a Case	
Süsen Banazılı and Mesut Öterkuş.....	<b>53</b>
Clinical experience in patients with Adult Still's disease	
Muhammed Okuyucu.....	<b>54</b>
Distal metatarsal Chevron osteotomy + modified McBride method in hallux valgus, short term 1-3 year results	
Muhammet Salih Ayas.....	<b>55</b>
Effects of Avanafil, A Phosphodiesterase Type 5 (Pde-5) Inhibitor, on A549 Lung Cancer Cell Proliferation	
Muhammet Çelik.....	<b>56</b>
Comparison of posterolateral approach and direct reduction posterior-anterior screw fixation method versus indirect reduction anterior-posterior screw fixation method for posterior malleolus fixation in ankle trimalleolar fractures	
Muhammet Kalkışım and Kerim Öner.....	<b>57</b>
Covid-19 and Comorbidities	
Mukadder Erdem, Recai Aci, Mahcube Çubukçu, Adem Keskin, Eda Türe and Ebru Ulaş.....	<b>58</b>
A Rare Form of Liver Failure and Immune Deficiency: IFNGR1 gene Mutation	
Murat Cag.....	<b>59</b>
Commitment to the medical profession and individual competitiveness of Medical Faculty students in Turkey	
Mustafa Bayraktar.....	<b>60</b>
Evaluation of staging 18F FDG PET/CT images with conventional and volumetric data in patients diagnosed with rectal cancer: Do the characteristics of the primary tumor provide information about the metastatic potential of the tumor?	
Nazlı Pınar Karahan Şen and Gamze Çapa Kaya.....	<b>61</b>

Is the Ganglion Cell Layer Thickness more sensitive marker than retinal nerve fibre layer thickness in Multiple Sclerosis Patients? Nuray Can Usta and Betül Önal Günay .....	62
How can the ocular surface cells be evaluated histologically with the impression cytology method? İrem İnanç, Burcu Kazancı, Fatma Çorak Eroğlu and Bizden Sabuncuoğlu .....	63
Does the tissue morphology change during the development of the age-related metabolic syndrome? Mihriban Alemdar, İrem İnanç, Deniz Billur and Nuray Yazıhan .....	64
Histological evaluation of the effectiveness of different microorganism types forming infections in the abscess formation Mehmet Batu Ertan, İrem İnanç, Mehmet Yağız Ayduğan, Ebru Evren, Esra Erdemli and Bülent Erdemli .....	65
Ultrastructural examination of anterior capsule samples taken from patients with congenital or juvenile cataract cases İrem İnanç, Pınar Bingöl Kızıltunç, Ferhad Özer, Huban Atilla and Belgin Can.....	66
The Psychosocial Effects of The Covid-19 Pandemic on Adolescents at A Private High School in Istanbul Gokce Hazar Otcu, Aydın Arman Canbaz, Sarp Esen and Meryem Merve Oren .....	67
Evaluation of the effect of mesenchymal stem cell administration on prognosis in critical COVID-19 patients in the intensive care unit Seyfi Kartal .....	68
Polymicrobial Sepsis Causes Edema and Neuronal Damage in Rat Cerebral Cortex; an Experimental Study Songül Doğanay and Özcan Budak .....	69
Pediatric Endocrinology and Syrian Refugee Children: A Photo We Want to Look Together Tuğba Kontbay and Müge Atar .....	70
Methanol Intoxication with Cerebral Hemorrhage: A Case Study Tuğçehan Sezer Akman, Hale Kefeli Çelik and Ahmet Şen .....	71
Is There a Relationship between Hip Fracture Surgical Treatment and Developing Dementia? One Year- Prospective Study Tuğrul Ergün .....	72
Effects of Virtual Reality on Stress Management of Disasters Volkan Ülker, Bedia Gülen and Yusuf Yürümez.....	73
Two siblings with Hyper IgE Syndrome diagnosed by Exome Sequence; DOCK8 mutation Yeşim Özdemir.....	74
Evaluation of protective effects of Ghrelin on Gastric tissue of ovariectomized rats Özlem Tuğçe Çilingir-Kaya.....	75
Effect of Chronic Alcohol Abuse on the Spectroscopically Determined Blood Secondary Structure of Proteins and Lipid Balance Zozan Güleken .....	76
Effects of Rosmarinic Acid Against Nephrotoxicity Induced by Cyclophosphamide in Rats Dilan Çetinavcı, Engin Yenilmez, Ayşe Firuze Bıyık, Ahmet Alver, Neslihan Sağlam.....	77
How Does Nonsteroidal Anti-Inflammatory Ibuprofen Affect Neural Tube Development in Chick Embryos? Esra Aslan and Hilal Güzel.....	78



# Lockdown effect on emergency surgical consults during the covid-19 outbreak: Experience of a pandemic hospital

Dr. Ahmet Burak Ciftçi<sup>1</sup> and Dr. Sönmez Ocak<sup>2</sup>

University of Health Sciences, Samsun Training and Research Hospital, Department of General Surgery, Samsun, Turkey <sup>1</sup>  
E-mail <sup>1</sup>: ahmetburak.ciftci@sbu.edu.tr, E-mail <sup>2</sup>: sonmezdr@gmail.com

## Abstract

**Background and aim:** The covid-19 pandemic has significantly impacted many lives and health care systems around the world. Many countries have imposed lockdowns and extraordinary restriction measures including curfews to control the disease. This study aimed to evaluate the lockdown effect on surgical emergencies during this outbreak.

**Methods:** Patients admitted to the emergency department (ED) and referred to general surgery (GS) was retrospectively recorded during the first lockdown (April 1, 2020-May 31, 2020) and before the covid-19 pandemic, corresponding time last year (April 1, 2019-May 31,2019). Patient admission rates to ED and general surgery outpatient clinics, patient characteristics, reasons for consultation, hospitalization rates, surgical intervention requirements, and reasons for urgent surgery were compared.

**Results:** In this study, a significant reduction was observed in the number of patients admitted to ED, and general surgery consultation is required in the covid-19 lockdown period (72% and 32% decrease respectively). There was no difference in hospitalization and surgical intervention rates between the groups ( $p=0.158$  and  $p=0.871$  respectively). The number of patients referred to general surgery with a diagnosis of perianal disorders was significantly higher in the lockdown group.

**Conclusions:** Covid-19 lockdown and restrictions have dramatically decreased general surgery patients' admissions to emergencies. Patients should be advised for future pandemics and lockdowns not to delay admissions to hospitals in emergent situations. We hope, this study findings will be guiding health authorities in making rearrangements in hospitals during subsequent waves and lockdowns in this pandemic.

**Keywords:** Covid-19 pandemic, lockdown, restrictions, emergency surgery

# Fusion Planet: Past, Today and Future of Computer Scientific Solutions in Digital Pathology

**Taha Yiğit Alkan, M.Sc.<sup>1</sup>, Asst. Prof. Hüseyin Gökhan Akçay<sup>2</sup> and Assoc. Prof. Havva Serap Toru<sup>3</sup>**

*Akdeniz University, Computer Engineering Department, Antalya, Turkey <sup>1,2</sup>*

*Akdeniz University, Department of Pathology, Antalya, Turkey <sup>3</sup>*

*E-mail <sup>1</sup>: yigitalkan@akdeniz.edu.tr, E-mail <sup>2</sup>: hgakçay@akdeniz.edu.tr, E-mail <sup>3</sup>: serapтору@akdeniz.edu.tr*

## Abstract

The increasing number and variety of the diagnostic and therapeutic techniques have led to a large data repository in pathology. Meanwhile, pathological data is usually processed with manual techniques. Although digitalization has started in the preanalytical period, it is not widespread and is open to further developments. Analytical and postanalytical periods need much more investigation which not only prevent time consumption but also provide laboratory and optical microscope free evaluation.

The commercial launch of Whole Slide Imaging (WSI) scanners has been a milestone in digital pathology (Pantanowitz et al., 2018). With the help of WSI scanners, it is possible to control the entire biopsy. Thus, the experience on the microscope can be transferred to digital platform which enables preservation of stains for re-evaluation and archiving.

Pathology is considered as gold standard for diagnoses in various medical applications such as oncology, surgery and transplantation where intra- and inter-observer disagreements in image-based decisions are frequently encountered (Allende et al., 2014; Nicholson, 2004; Pournik et al., 2014). However, mathematical modeling of pathological images can lead to the standardization of these diagnoses (Choi et al., 2020). Through these models, machines are able to learn from past data to create classification and scoring tools that can assist pathologists in making revisable and reviewable decisions. A machine learning decision system usually requires domain expertise to transform the raw image pixels into a suitable feature vector for representing decisions of interest. Unlike traditional approaches, deep learning does not need hand-crafted image features that are possibly not well-suited to the problem. Rather, deep neural networks take example images and expected decisions as input and automatically learns hidden image features most relevant to the application. Hence, these automatic methods become feasible also for non-experts.

Although diagnostic AI studies on cancer/metastasis detection, scoring, grading, mitosis, and proliferation index counting have started in the early 1990s (Moxley-Wyles et al., 2020), the first certificated AI algorithm for pathology was developed in 2018 (Pantanowitz et al., 2020). While all healthcare practices are undergoing digital transformation, pathology is still mostly analogous. Augmentation of AI technology in the fields of immunohistochemistry, special histochemistry, immunofluorescent staining, and molecular pathology can decrease the workload, provide label-free evaluation and preserve tissue for molecular techniques. Although there are many challenges and doubts, there is still hope for reaching molecular data from morphology. Providing technical infrastructure for molecular examinations is not very easy from technical and economical perspectives on routine applications. However, in cases where digital pathology infrastructure is established, accessing molecular information from morphology with AI algorithms can be a cost-effective solution to be able to perform applications in every medical center. Current practices and research show that pathologists will play an important role in multidisciplinary and interdisciplinary



studies (Rashidi et al., 2019). Even though the data is sufficient and of high quality, computer science will be indispensable in the production of automated solutions for pathology. Also, standardization of these practices will cause legal, ethical, and reliability problems that will reveal new challenges.

**Keywords:** *Digital Pathology, Artificial Intelligence, Image Processing*

## 14q32.31q32.33 Microdeletion Detected by Chromosomal Microarray in a Child with Dysmorphism and Hypotonia

Aslihan Sanrı, MD<sup>1</sup> and Özlem Sezer, MD<sup>2</sup>

*Department of Pediatric Genetics<sup>1</sup>, Department of Medical Genetics<sup>2</sup>, University of Health Sciences, Samsun Training and Research Hospital, Samsun, Turkey  
E-mail: aslihansanrı@yahoo.com*

### Abstract

Terminal and interstitial deletion of chromosome 14q32 region are rare. Although this is quite rare, the phenotype of 14q32 deletion syndrome has been described before. Common clinical features are prenatal and/or postnatal growth deficiency, developmental delay, intellectual disability, microcephaly, hypotonia, high forehead, broad and flat nasal bridge, blepharophimosis, ptosis, epicanthus, short bulbous nose, long and broad philtrum, thin upper lip, high arched palate, abnormal dentition, malformed helices, low set ears, short neck, single palmar crease, and congenital heart defects. We report a patient with interstitial 14q32.31q32.33 deletion detected by chromosomal microarray. A two-month-old girl was referred to genetic department because of facial dysmorphism and hypotonia. Parents were nonconsanguineous and the family history was unremarkable for intellectual disability, developmental delay, and congenital malformations. She was born at term with a birth weight of 2960 g (10-50th percentile). On physical examination, her weight was 3330 g (<3rd percentile), height was 54 cm (3-10th percentile) and head circumference was 36 cm (<3rd percentile). Facial dysmorphic features included hypertelorism, epicanthus, short nose, high forehead, deep and long philtrum, thin upper lip, retro micrognathia, high arched palate and short neck. She had mild hypotonia and she could not hold her head. Kidney and trans fontanelle ultrasonography scans were normal. Echocardiogram revealed a secundum type atrial septal defect. Hearing spelling and ophthalmological examinations were normal. Because of postnatal growth deficiency, dysmorphism, hypotonia and congenital cardiac anomaly chromosome analysis and microarray were requested. The patient's chromosome analysis was normal, 46, XX. Microarray showed a 3.7 Mb deletion at chromosome 14q32.31q32.33. A total of 72 genes were deleted and 16 of them are associated with known phenotypes. Parental chromosome analysis and microarray were normal and showed de novo origin of the deletion. Among the patients with 14q32 microdeletions, the majority had terminal deletions, in comparison to interstitial deletions. Only a few cases of interstitial 14q32 microdeletion reported in current literature to date. Our case had interstitial deletion and there is any reported case with the similar breakpoints with our patient. We believe that our case contributes to additional phenotype seen in patients with interstitial 14q32 deletion. With increasing frequency of patients with dysmorphic features, developmental delay and intellectual disability are found to have a chromosomal imbalance which is submicroscopic often and can be detected with the specific cytogenetic techniques such as chromosomal microarray. The advances in the use of microarrays have permitted the increasing number of reports of several microdeletion and microduplication syndromes especially in patients with developmental delay and/or intellectual disability. Chromosomal microarray also allows for a more detailed description of location, size and genes involved in a specific chromosome region, and are helpful to characterize the genotype-phenotype correlations more precisely.

**Keywords:** *The globalization of science; scientific collaboration; globalizing knowledge economy*

# Factors Associated with Pressure Ulcers in Adult Critical Care Patients

**Dr. Başar Erdivanlı**

*Recep Tayyip Erdogan University, Medical Faculty, Rize, Turkey  
E-mail: basar.erdivanli@erdogan.edu.tr*

## Abstract

**Background:** Pressure ulcers represent a significant problem in critical care patients. Presence of ulcers increase duration of hospital stay, morbidity and complexity of the patient care. It is known that hospital-acquired pressure ulcers are multifactorial. Due to the low number of randomized controlled studies, systematic reviews reported several factors, of which age, diabetes mellitus, cardiovascular disease, hypotension, prolonged hospital stay and mechanical ventilation, and vasopressor administration appeared in all reviews.

**Purpose:** This study investigated the potential of a relational database to predict occurrence of pressure ulcers.

**Methodology:** An adult intensive care database was queried to find factors related to occurrence, worsening, or healing of pressure ulcers. No laboratory data was used.

**Results:** The data from a total of 611 patients were evaluated. Patients with pressure ulcer at the day of admission to critical care unit (n = 38) were excluded from the final analysis. It was understood that all patients received the same skin care, repositioning and support surfaces. The level of education of the staff was similar throughout the period of data collection.

Pressure ulcers were observed in 92 patients (16%). Major determinants of development of pressure ulcers were admission ApacheII and SOFA scores ( $p < 0.002$  and  $p = 0.025$ , respectively). Specific characteristics such as age, body proportions, unintentional weight loss and presence of diabetes mellitus or hypertension, and diagnosis at the admission were not predictive for development of pressure ulcers. Patients who had pressure ulcers were slightly older ( $69 \pm 15$  vs  $64 \pm 19$  years,  $p = 0.028$ ), longer critical care unit stay ( $15-32$  vs  $1-8$  days,  $p < 0.001$ ), and were more commonly discharged to palliative care ( $30\%$  vs  $5\%$ ,  $p < 0.001$ ). Propensity score matching according to development of pressure ulcers yielded similar results.

**Conclusion:** This retrospective study showed that a relational database may predict development of pressure ulcers by commonly used metrics like ApacheII and SOFA scores. The inclusion of several factors in these scoring systems support the notion that occurrence of pressure ulcers cannot be explained by a single factor. This may explain the neutral effect of factors such as sedation or vasopressors. We are in opinion that skin care, repositioning and use of support surfaces played a role in the low occurrence of pressure ulcers.

**Keywords:** *Pressure ulcer, Critical Care, Risk scores, Adult, Human.*

# COVID-19 Through the Eyes of Psychiatrists: Back Pain Can be a Symptom and May Predict Pneumonia in COVID-19

**Dr. Cuma Uz**

*Kırıkkale High Specialized Hospital, Physical Medicine and Rehabilitation Clinic, Kırıkkale, Turkey  
E-mail: cumauz12@gmail.com*

## Abstract

**Objective:** The aim of the study was to determine whether back pain is a clinical manifestation in patients with COVID-19, and to determine whether any demographic and disease characteristics could act as an effective indicator of back pain.

**Design:** The patients with COVID-19 (N: 99) were recruited from the infectious diseases department of the secondary care hospital and *divided into two groups according* to the presence or absence of back pain. The main outcomes included were demographic and disease characteristics, Nord-Trøndelag Health Study Physical Activity Level for Work (HUNT), 6-minute walking test (6MWT).

**Results:** The most common symptoms were fatigue (n = 63, 63.6%), followed by back pain (n = 50, 50.5%). Sedentary lifestyle, oxygen requirement, presence of pneumonia and typical pneumonia pattern were significantly higher (p = 0.009, p = 0.026, p = 0.001, p = 0.001, respectively), and aerobic capacity was lower (p = 0.001) in the patients with back pain. The presence of back pain continued to be associated with the presence of pneumonia in multivariate analysis.

**Conclusions:** Back pain may be associated with the presence of COVID-19 pneumonia and should be evaluated as an early warning symptom.

**Keywords:** *Coronavirus disease 2019 (COVID-19), SARS-CoV-2, back pain, pneumonia*

# The Effect of Ulipristal Acetate and Vitamin D3 on Folliculogenesis

**Dr. Damla Gül Fındık<sup>1</sup>, Dr. Gülnur Take Kaplanoğlu<sup>2</sup>, Gökçe Nur Arık, M.Sc.<sup>3</sup>  
and Dr. Nagya B. Abubaker Alemari<sup>4</sup>**

*Gazi University, Faculty of Medicine, Department of Histology and Embryology, Ankara, Turkey*<sup>1,2,3,4</sup>  
*E-mail<sup>1</sup>: damla.findik@gmail.com, E-mail<sup>2</sup>: gulturtake@gmail.com, E-mail<sup>3</sup>: gnurarik@gmail.com,  
E-mail<sup>4</sup>: nagwa.alamari@gmail.com*

## Abstract

Folliculogenesis is a complex process that occurs in the ovaries, with various growth factors and signal molecules take part. One of the hormones involved in the control of ovulation is progesterone. Recent studies show that progesterone stimulates ovulation and primary follicle development. High progesterone concentration suppresses ovulation and secondary follicle development. In the light of this information, it can be deduced that Ulipristal acetate (UPA) as a progesterone receptor modulator has a role in folliculogenesis. There are limited studies about the effect of UPA on folliculogenesis. Vitamin D3 is another factor that has been shown to play an important role in ovarian functions including follicular development. VitD3 supplementation increases preantral follicle survival, antral follicle growth, and survival. It has been shown that VitD3 also regulates steroidogenesis by stimulating progesterone and estradiol synthesis in the ovary. Research on follicular development mechanisms will provide more comprehensive data on female reproductive life and the development of new therapeutic approaches against reproductive aging. In this respect, it is important to understand the effect mechanism in folliculogenesis of UPA and VitD3, which are increasingly used together in recent years for antifibroid effects in uterine fibroid treatment. In our study, we investigated the effects of UPA and VitD3 on folliculogenesis by used histochemical methods in a rat model. In study 48 female Wistar-albino rats randomly divided into seven groups: control group, 3 weeks oral VitD3 (1000 IU/kg/day) administration group, 5 weeks oral UPA (3 mg/kg/day) administration group, every 2 days for 5 weeks oral DES (1,35 mg/kg/day) and IM progesterone (1mg/kg) administration group, after DES and progesterone administration 3 weeks oral VitD3 (1000 IU/kg/day) treatment group, 5 weeks oral UPA (3 mg/kg/day) treatment group, VitD3+UPA treatment group. Follicles in normal stages have become atretic and hemorrhagic cystic structures were observed in the UPA group. In the high estrogen–progesterone administration group, the granulosa cell layers of antral follicles were determined to be relatively thin like the polycystic ovary syndrome. Giant follicles with thinning walls were also observed in the UPA and VitD3+UPA treatment groups. In the high estrogen–progesterone applied group, many corpus luteums were detected due to an increase at LH peak and ovulation caused by estrogen. Corpus luteum degeneration was also observed before maturation because of negative inhibition of progesterone. The structure of the corpus luteum was normal in the VitD3 treatment group. We demonstrated that; UPA prevented progesterone negative feedback, and corpus luteums were observed to be enlarged in the UPA treatment group. UPA and VitD3 cause excessive vascular dilatation and congestion. As a result, it was observed that VitD3 and VitD3+UPA treatment groups gave the best results in terms of folliculogenesis in the condition of high estrogen – progesterone.

**Keywords:** *Folliculogenesis; Progesterone; Ulipristal acetate; Vitamin D*

## Evaluation of Antibody Levels in Pcr Negative Covid-19 Suspected Case Series

**Asst. Prof. Demet Gür Vural<sup>1</sup>, Res. Asst. Büşra Usta<sup>2</sup> Assoc. Prof. Yeliz Tanrıverdi Çaycı<sup>3</sup>, Asst. Prof. Kemal Bilgin<sup>4</sup> and Prof. Asuman Birinci<sup>5</sup>**

*Institution: Ondokuz Mayıs University, Department of Medical Microbiology, Faculty of Medicine, Samsun, Turkey<sup>1</sup>  
E-mail<sup>1</sup>: demet.gur@yandex.com , E-mail<sup>2</sup>: dr.busraozkaya@gmail.com, E-mail<sup>3</sup>: yeliztanriverdi@gmail.com, E-mail<sup>4</sup>: kemal.bilgin@omu.edu.tr, E-mail<sup>5</sup>: asumanbirinci@yahoo.com*

### Abstract

**Introduction:** Real Time-Polimerase Chain Reaction (RT-PCR) test is used as the gold standard test in the diagnosis of COVID-19. Mostly, viral RNA becomes detectable in-patient samples up to 3 days before the onset of disease symptoms with the PCR method. The sensitivity of SARS-COV2 RNA tests is reported to be 55-75 percent. A negative PCR result is not sufficient to exclude the disease in the presence of compatible symptoms and imaging findings. These patients are evaluated as suspected COVID-19 and treated.

Examination of antibodies against the SARS-COV2 virus in patients whose PCR test is negative from patients with suspected COVID-19 disease may be helpful in diagnosis. It has been reported that SARS-COV2 specific antibodies are formed within 5-14 days from the onset of disease symptoms. In our study, we investigated SARS-COV2 Ig G levels in patients with negative RT PCR and clinically suspected Covid-19.

**Material Methods:** The data of 9 patients whose symptoms were compatible with COVID-19 and whose RT-PCR test was negative were examined from the data of patients between August-October 2020 at Ondokuz Mayıs University Hospital. Two months after the negative test results, patients were called, and venous blood samples were collected. SARS-COV2 Ig G was investigated by Enzyme Linked Immunoassay (ELISA) method according with the manufacturer's recommendations (Euroimmun, Germany).

**Result:** The age range of the patients was between 29-72 and 6 patients (66.6%) had comorbid disease. Pulmonary computed tomography was performed in 8 of 9 patients and the results of 6 patients (75.0%) were reported in accordance with COVID-19. Eight of 9 were hospitalized and 1 patient was taken to the intensive care unit. SARS-COV2 Ig G was found to be positive in 8 patients (88.8%) in blood samples taken 2 months later.

**Conclusion:** There may be many reasons behind this negativity in the PCR results, such as the symptom day on which the swab samples were taken, the way they were taken, the transport or RNA extraction stage, and the sensitivity of the kits used in the study.

The use of antibody tests against SARS-COV2 together with RT-PCR in the diagnosis of suspected patients may be useful in the diagnosis of patients.

**Keywords:** RT-PCR, SARS-COV2 Ig G, Covid-19

# Evaluation of Antinuclear Antibodies in Patients who were Infected with Covid 19

**Asst. Prof. Demet Gür Vural**

*Ondokuz Mayıs University, Department of Medical Microbiology, Faculty of Medicine, Samsun, Turkey  
E-mail: demet.gur@yandex.com*

## Abstract

**Introduction:** Coronavirus Disease 2019 (COVID-19), caused by Severe Acute Respiratory Syndrome Coronavirus-2 (SARS-CoV-2) infection, is associated with many different clinical features that are commonly found in autoimmune diseases, including arthralgias, myalgias, fatigue, sicca, and rashes. Less common manifestations of autoimmunity have also been observed in COVID-19 patients, including thrombosis, myositis, myocarditis, arthritis, encephalitis, and vasculitis. These clinical observations, and the increasing proportion of “recovered” patients with persistent post-COVID-19 symptoms suggest that inflammation in response to SARS-CoV-2 infection promotes tissue damage in the acute phase and potentially some of the long-term sequelae. In our study, we aimed to evaluate the frequency of Antinuclear Antibodies in patients after Covid-19 infection.

**Material Metods:** 45 patients who had infected Covid -19 and ANA test has been studied were included in our study. ANA IIFA results were retrospectively evaluated. The presence of ANA and staining pattern was evaluated with Indirect Immunofluorescence Antibody (IIFA). The commercial IIFA kit (Euroimmun AG, Lübeck, Germany), which contains HEP-2 and monkey liver cells together as a tissue for ANA IIFA testing, was used. Prepared preparations were evaluated at a fluorescence microscope (Euroimmun AG, Lübeck, Germany) at a magnification of 400x. The results were reported qualitatively (+, ++, +++, +++) according to the fluorescence intensity of the slides and their patterns.

**Results:** Of the patients included in the study, 29 (64,5%) were female and 16 (35,5%) were male. When we look at the distribution of ANA patterns in positive samples; speckled 10 (22,2%), speckled and cytoplasmic granular 6 (13,3%), homogen and speckled 8(17,8%), granuler and speckled 6 (13,3%) were the most common (Table 1).

**Table 1. Distribution of ANA patterns**

ANA Patterns	%
Speckled	10(22,2%)
Homogen+speckled	8(17,8%)
Speckled+nucleoler	6(13,3%)
Speckled+granular in thecytoplasm	6(13,3%)
Speckled+ Granularchromosomes	2(4,5%)
Nucleoler+homogene	1(2,2%)
Nucleoler+ granular in thecytoplasm	2(4,5%)
Speckled in the cytoplasm	4(9%)
Nucleoler	3(6,6%)
Mitotic cell	3(6,6%)

**Conclusion:** Though the exact etiology of autoimmune diseases still remains unknown, there are various factors which are believed to contribute to the emergence of an autoimmune disease in a host

including the genetic predisposition, the environmental triggers such as bacterial infections, including the gut microbiota, viral fungal and parasitic infections, as well as physical and environmental agents, hormonal factors and the hosts immune system dysregulation. In order to determine the relationship between Covid 19 and autoimmune disease, there is a need for studies with extensive patient groups and the follow-up of autoimmune markers before and after the disease.

**Keywords:** *Covid 19, Anti-nuclear antibody, autoimmunity*



# Depression, Hopelessness, and Social Support in Hemodialysis Patients

**Dr. Demet Yavuz**

*Department of Internal Medicine, Division of Nephrology, Samsun Training and Research Hospital, Samsun, Turkey  
E-mail: demetdolu@hotmail.com*

## Abstract

**Introduction:** The hemodialysis regimen is an inevitable and mandatory treatment for patients with end-stage renal disease (ESRD). During the dialysis journey, patients may experience depressive symptoms, and hopelessness. These problems are common in hemodialysis patients. Social support has been consistently linked to better health outcomes in this group patients. The objective of the present study was to investigate the link between depression, hopelessness, and social support in hemodialysis patients.

**Materials and Methods:** A total of 131 patients were included in study (58 woman; mean age  $48.9 \pm 14.2$  years; hemodialysis duration  $40.6 \pm 25.9$  months). Demographic data and laboratory values were evaluated. We used Beck Depression Inventory (BDI) and Multidimensional scale of perceived social support (MSPSS) and Beck Hopelessness Scale (BHS) in all patients.

**Results:** BDI, MSPSS and BHS were  $14.1 \pm 6.3$ ,  $47.6 \pm 19.8$ ,  $5.6 \pm 3.3$  respectively. There was negative correlation between BDI and MSPSS ( $r = -0.485$ ,  $p = 0.001$ ), and BHS inversely correlated with MSPSS ( $r = -0.560$ ,  $p = 0.001$ ). In the multivariate linear regression analysis, BDI was independently associated with MSPSS ( $\beta = -0.154$ ; 95% confidence interval,  $-0.204$  to  $-0.104$ ,  $p < 0.001$ ), BDI was independently associated with female gender ( $\beta = 5.503$ , 95% confidence interval  $3.715$  to  $7.291$   $p < 0.001$ ) and single person ( $\beta = 2.815$ , 95% confidence interval  $0.868$ - $4.761$   $p = 0.005$ ). In the multivariate linear regression analysis, BHS was independently associated with MSPSS ( $\beta = -0.031$ ; 95% confidence interval,  $-0.055$  to  $-0.007$ ,  $p < 0.05$ ), BHS was independently associated with female gender ( $\beta = 1.542$ , 95% confidence interval  $0.793$  to  $2.290$   $p < 0.001$ ) and single person ( $\beta = 0.759$ , 95% confidence interval  $0.029$ - $1.489$   $p < 0.05$ ).

**Conclusion:** The result of this study indicates that DBI and BHS are negatively correlated with MSPSS. Hemodialysis patients needed more social and psychological support. The lower social support that associated with the presence of depression and hopelessness.

**Keywords:** *Depression, Hopelessness, Social Support, Hemodialysis*

# Effects of thymoquinone on anxiety-related behavior and auditory potentials in rats exposed to glucocorticoid

**Dr. Deniz Kantar**

*Department of Biophysics, Faculty of Medicine, Akdeniz University, Antalya, Turkey  
E-mail: dkantar@akdeniz.edu.tr*

## **Abstract**

The present study was carried out to determine the role of thymoquinone (TMQ) in modulating the auditory evoked responses, anxiety-related behavior, and serotonin level in rats that stress induced by corticosterone (cort) administration. Rats were divided into 4 groups (n=6); Sham(S), TMQ (TMQ), corticosterone (CORT) and Cort+TMQ (CTMQ) groups. Corticosterone (3 mg/kg) in normal saline was administered intraperitoneally (i.p.) and TMQ (20 mg/kg) in corn oil was administered by gavage. After half an hour, elevated plus maze test were performed to assess anxiety-related behavior. Besides, loadness-dependent auditory evoked potentials were recorded and loadness dependence and oscillatory responses were analyzed. Then, rats were sacrificed and their brains were removed for serotonin estimation. Cort injection resulted in a reduction in the open arm duration ratio in the EPM test. TMQ treatment prevented the observed alterations in the CTMQ group. Cort injection led to a significant increase in the levels of loadness dependence of auditory potentials and decreased auditory delta/theta responses. TMQ treatment reversed the cort related alterations in auditory evoked potential parameters. Moreover, cort injection resulted in a reduction in the serotonin content and TMQ treatment prevented the observed alterations in the CTMQ group. In conclusion, this study demonstrates that TMQ prevented cort-induced serotonergic changes, which may partly contribute to the improvement of auditory network dynamics and anxiety-related behavior in rat.

**Keywords:** *Thymoquinone, Anxiety, Auditory Evoked Potential, Serotonin*

# Spontaneous Tumor Lysis Syndrome in Lung Cancer: Very Rare Case for Solid Tumors

Dr. Buğra Özel<sup>1</sup>, Dr. Düriye Sıla Karagöz Özen<sup>2</sup> and Dr. Ahmet Baran<sup>3</sup>

Health Sciences University, Samsun Research and Training Hospital,  
Department of Internal Medicine <sup>1,2</sup>, Department of Medical Oncology <sup>3</sup>  
E-mail <sup>1</sup>: bbugrab@gmail.com, E-mail <sup>2</sup>: silakaragoz@yahoo.com, E-mail <sup>3</sup>: drahmetbaran19@gmail.com

## Abstract

**Introduction/purpose:** Here we report a 67-years-old male patient with a diagnosis of lung cancer who presented with spontaneous tumor lysis syndrome.

**Case:** A 67-year-old male patient was admitted to the emergency department with physical instability, weakness, and fatigue. He has diagnosed with neuroendocrine lung cancer 1 year ago. He was treated with topotecan chemotherapy 21 days before admission. Physical examination revealed conjunctival pallor, scleral icterus. Cardiovascular system and respiratory system examinations were normal. Tender hepatomegaly was palpated. The patient was oliguric. Laboratory results were summarized in table 1. Twenty-one days after the last chemotherapy, the patient was diagnosed with spontaneous tumor lysis syndrome as a result of clinical and laboratory examinations and hospitalized. Hydration was started rapidly with intravenous NaCl 0.09% with the diagnosis of acute renal failure and tumor lysis syndrome. Oral and intravenous hydration treatments were done according to their close follow-up. Venous blood gas, urea, creatine, and electrolyte follow-ups were performed daily. There was no need for hemodialysis during the patient's hospitalization. Hyperuricemia, hyperkalemia, and hyperphosphatemia resolved. He was discharged on the seventh day of hospitalization. The last laboratory findings are summarized in Table 1.

**Conclusion:** This is a tumor lysis syndrome case in a patient with a solid tumor diagnosis. Although it can be observed during hematological malignancies, spontaneous tumor lysis syndrome is a very rare entity for solid tumors. This report is important for clinicians to keep in mind.

**Table 1.** Laboratory results

	First Visit	Last Visit	Referans Range	Units
pH	7.297	7.36	(7.35-7.45)	
PaCO <sub>2</sub>	39.4	32	(35.0-45.0)	mMol/L
HCO <sub>3</sub>	19.2	18.3	(22.0-24.0)	mMol/L
K <sup>+</sup>	5.61	3.69	(3.5-5.1)	mEq/L
Ca <sup>+2</sup>	9.65	8.79	(8.6-10.2)	mg/dl
P	5.05	3.59	(2.8-4.5)	mg/dl
Uric asid	19.76	6.67	(3.4-7.0)	g/dl
Creatine	2.94	1.11	(0.6-1.2)	mg/dl

**Keywords:** Tumor Lysis Syndrome, Lung Cancer, Solid Tumors

## Is Selenium deficiency related to an increased risk for diabetic retinopathy?

Dr. Hacer Pınar Öztürk Kurt<sup>1</sup>, Dr. Düriye Sıla Karagöz Özen<sup>2</sup>, Dr. İpek Genç<sup>3</sup>,  
Dr. Mukadder Erdem<sup>4</sup> and Prof. Mehmet Derya Demirag<sup>5</sup>

*Clinic of Internal Medicine, Health Sciences University, Samsun Education and Research Hospital, Samsun, Turkey*<sup>1,2,5</sup>

*Clinic of Ophthalmology, Health Sciences University, Samsun Education and Research Hospital, Samsun, Turkey*<sup>3</sup>

*Clinic of Biochemistry, Health Sciences University, Samsun Education and Research Hospital, Samsun, Turkey*<sup>4</sup>

*E-mail*<sup>1</sup>: hpinarim@hotmail.com, *E-mail*<sup>2</sup>: silakaragoz@yahoo.com, *E-mail*<sup>3</sup>: albayrakipek@yahoo.com,

*E-mail*<sup>4</sup>: dr.mukadder@hotmail.com, *E-mail*<sup>5</sup>: mehmetderyademirag@yahoo.com

### Abstract

**Aim:** Diabetic retinopathy is one of the leading causes of visual loss among adults. Oxidative stress plays an important role in the pathogenesis of diabetic retinopathy. This study aims to determine whether selenium deficiency is related to the elevated risk of diabetic retinopathy in patients with type 2 diabetes mellitus.

**Materials and Methods:** The patients were selected among patients who applied to the Health Sciences University, Samsun Research and Training Hospital Internal Medicine outpatient clinics. 115 Patients with type 2 Diabetes Mellitus (DM) were included in the study. The retinopathy group included 47 patients, and the non-retinopathy group included 68 patients. Plasma samples were collected from the patients to determine selenium levels.

**Results:** Mean age of the retinopathy group was 56.5±10 years and the mean age of the non-retinopathy group was 53.2±9 (p = 0.070). Gender distribution between the two groups was similar (p = 0.801). The mean selenium level of the retinopathy group was 70.11±17.28 µg/l, and the mean selenium level of the non-retinopathy group was 80.20±19.10 µg/l. The mean selenium level of the retinopathy group was significantly lower than that of the non-retinopathy group (p = 0.005). The median duration of DM was significantly higher in the retinopathy group than that of the non-retinopathy group [10 (1-25) and 6 (1-21) respectively and p = 0.002]. Logistic regression analyses showed that higher levels of blood selenium values were an independent preventive factor against retinopathy occurrence [OR and 95% CI=0.965(0.939-0.991)] while the duration of DM was an independent risk factor for retinopathy occurrence [OR and 95% CI= 1.131 (1.050-1.219)].

**Conclusion:** In our study, selenium levels differed significantly between the retinopathy group, and the non-retinopathy group. Our findings may improve preventive choices against diabetic retinopathy. Besides controlling hyperglycemia and high blood pressure, we can measure blood selenium levels of patients with diabetic retinopathy and replace it if deficient.

**Keywords:** *Diabetes mellitus, diabetic retinopathy, oxidative stress, selenium deficiency*

# The Development of Family Medicine Identity Scale

Dr. Duygu Üstünol<sup>1</sup>, Dr. İsmail Kasım<sup>2</sup> and Prof. Dr. Adem Özkara<sup>3</sup>

*Sulakyurt State Hospital, Kırıkkale, Turkey<sup>1</sup>*

*Health Sciences University, Ankara City Hospital, Ankara, Turkey<sup>2,3</sup>*

*E-mail<sup>1</sup>: duyguustunol@gmail.com, E-mail<sup>2</sup>: astronomkasim@yahoo.com, E-mail<sup>3</sup>: ademozkara@yahoo.com*

## Abstract

**Objective:** It is to put a scale into use of the academic community which measures how well the doctors can interiorize the features concerning family practice the training of family practice specialization in our country, by developing a family practice identity scale that is peculiar to Turkey.

**Materials and Methods:** It is the study of developing a scale. 5-point Likert scale is generated by creating the questions about core proficiencies of family medicine were defined by WONCA and professional identity. The scale is evaluated for extent and content by a 16-person specialist group. The aforementioned scale is studied to verify the validation and reliability in Turkey. Bartlett's test result is analyzed to be determined the situation of factor analysis availability of data. The subscales of the scale are described by exploratory factor analysis. Cronbach's Alfa value is estimated to be specified item compliance values of obtained factors. In addition to this, Mann Whitney U test, ANOVA test, Kruskal Wallis test are performed where necessary. The scale has been performed on 351 people who work as academicians, specialists, and residents in family medicine in Turkey between 23.05.2019-07.07.2019. In order to expedite the interpretability of acquired scores, the subscale scores and total scores are transformed according to the 100-point system. To convert, the below-mentioned formula is utilized.

$$Score_{100} \left( \frac{\text{Obtained total score}}{\text{Obtained maximum score from the subscale}} \right) * 100$$

**Findings:** The answers given by 350 doctors are reported according to the statistical analyses performed. Of the 350 family physicians who participated in the study, %64,6 (n=226) are women and %35,4 (n=124) are men. Of the participant, %57,1 (n=200) are family medicine residents, %29,7 (n=104) are family medicine specialists, %13,1 (n=46) are family medicine academicians. Participants' median age is appointed as 31.00 and their mean age as 34.72. The average age of the female participants is 32, and the median age is 29. The average age of the male participants is as 39, and their median age is 35. During the analyses, while the first four factors whose eigenvalues are the highest are kept fixed, the questions from the other factors are distributed according to their content similarities. As a result, the scale of four-factor structure with forty-six questions is obtained. The sub-scales are named by the contents of the questions: Patient-doctor communication, professional satisfaction, the scope of the working area and comprehensive approach, biopsychosocial approach. It has been concluded that the scale is a valid and reliable questionnaire in Turkey after these advanced statistical analyses.

**Discussions and Result:** "The Scale of Family Practice Identification" is developed successfully. With the aforementioned scale, by observing professional progress of residents, the doctors that have occupational identity and sense of belonging can be trained for the community of family practice.

**Keywords:** *Family medicine, family medicine identity, professional identity, scale development*

# Investigation of the relationship between magnesium level and vitamin D, bone mineral density, knee osteoarthritis and chronic diseases

Dr. Ebru Yilmaz<sup>1</sup> and Dr. Sena Ünver<sup>2</sup>

Kocaeli Government Hospital, Department of Physical Medicine and Rehabilitation, Kocaeli, Turkey<sup>1,2</sup>  
E-mail<sup>1</sup>: dr.ozcanebru@gmail.com, E-mail<sup>2</sup>: akcasena83@gmail.com

## Abstract

Magnesium (Mg) is obligatory for maintaining numerous physiological cellular functions. Mg deficiency is linked with a number of health conditions including osteoporosis, hypertension, diabetes mellitus, atherosclerosis and coronary heart disease, and malignancies (colon and breast). Although calcium (Ca) and vitamin D have been the master focus of nutritional prevention of osteoporosis, several minerals such as copper, zinc, selenium, and Mg are also known to be important. Mg is dominantly located within the cartilage and bone of a human body. In Mg deficiency, there are decreased synthesis, release, and action of parathyroid hormone (PTH) and 1,25(OH)<sub>2</sub>D. In several studies, a significant association has been found between bone density and the intake of Mg and dietary Mg restriction promotes osteoporosis. Moreover, there is some evidence about the link between Mg level and prevalence of knee osteoarthritis (OA). The aim of the study was to evaluate the relationship between Mg level and vitamin D<sub>3</sub>, bone mineral densitometry (BMD), knee OA and chronic diseases. A total of 98 patients (92 female, 6 male) between the ages of 40 and 75 who presented to the outpatient clinic with complaints of knee pain were included. Data on age, sex, body mass index (BMI), smoking, menopausal status, duration of menopause, family history of osteoporosis, the presence of chronic diseases (hypertension=HT, diabetes mellitus=DM, hyperlipidemia=HPL, coronary artery disease=CAD, hypothyroidism=HPT). The serum levels of Mg, Ca, 25(OH)-vitamin D<sub>3</sub>, alkaline phosphatase (ALP) and PTH measurements were performed to whole patients. Moreover, all patients underwent weight-bearing bilateral anteroposterior radiography of the knee by using X-Ray, and BMD of femoral neck and lumbar vertebrae (L1-L4) by using dual-energy X-Ray absorptiometry (DEXA). The presence of osteoporosis was accepted as T scores  $\leq -2.5$ . The mean age of the study population was  $59.15 \pm 10.58$  years. Forty-seven (48%) patients had osteoporosis: two of them was male (4.3%) and the others were female (95.7%). Of all patients, 8.2% (n=8), 22.4% (n=22), 45.9% (n=45), 20.4% (n=20), and 3.1% (n=3) had KL Grade 0, 1, 2, 3, and 4 knee OA, respectively. The percentages of chronic diseases were 35.7% for HT (n=35), 20.4% for DM (n=20), 4.1% for HPL (n=4), 10.2% for CAD (n=10), and 21.4% for HPT (n=21). A statistically significant relationship was found between the level of Mg and age, smoking, duration of menopause, presence of chronic disease, PTH level, vitamin D level and femoral neck T score ( $p < 0.05$ ). The optimizing Mg status through diet and supplementation seems to be a safe and beneficial treatment for the regulation of vitamin D and PTH metabolism, osteoporosis and various chronic diseases. Future studies are needed to investigate the relationship between Mg and knee OA.

**Keywords:** Magnesium; vitamin D; osteoporosis; chronic diseases; knee osteoarthritis

# Outcomes of Breast Cancer Brain Metastasis Patients Undergoing Stereotactic Radiotherapy

Dr. Ela Delikgöz Soykut<sup>1</sup> and Dr. Nilgün Şahin<sup>2</sup>

Health Sciences University, Samsun Training and Research Hospital, Radiation Oncology Clinic, Samsun, Turkey <sup>1</sup>

Health Sciences University, Samsun Training and Research Hospital, Radiation Oncology Clinic, Samsun, Turkey <sup>2</sup>

E-mail <sup>1</sup>: eladelikgoz@gmail.com, E-mail <sup>2</sup>: nilgun\_orkun@hotmail.com

## Abstract

**Background:** In recent years, stereotactic radiotherapy (SRT) has been preferred to whole brain radiotherapy (WBRT) in the treatment of patients diagnosed with brain metastasis, especially considering the number and size of the lesions.

**Purpose:** It was aimed to examine the treatment results of patients who underwent SRT with a diagnosis of breast cancer brain metastasis.

**Methodology:** Between January 2015 and September 2018, breast cancer brain metastasis patients who received SRT alone or were administered SRT and WBRT together, or those who underwent SRT due to progression were retrospectively analyzed. The clinical and pathological characteristics of the patients were accessed from the automation system and the patient file archive. Survival curves were determined by Kaplan-Meier analysis, and prognostic significance of variables showing significance in univariate analysis were determined by multivariate analysis.

**Results:** 45 patients were included in the study. Median age was 51 (19-83), median follow-up time was 20 (1-70) months. Twenty-two (36.1%) of the patients diagnosed with breast cancer were stage 4 at the time of diagnosis, and 2 (3.3%) patients had brain metastasis at the time of diagnosis. There were single brain metastasis in 14 (31.1%) patients, 2-5 in 15 (33.3%), and multiple brain metastasis in 16 (35.5%) patients. SRT alone was applied to 17 of the patients (37.7%), SRT and WBRT was applied to 11 (24.4%) of the patients, and WBRT alone was applied to 17 (37.7%) of the patients. SRT doses ranged from 15-24 Gy / 1 frx, 9 Gy / 2 frx, 6-8 Gy / 3 frx. Median overall survival (OS) for all groups was 26 (95% CI, 17,1-34,8) months, 1-y, 3-y and 5-y OS were 75.6%, 40.5% and 24.6%, respectively. In terms of radiotherapy type, SRT alone, WBRT alone or WBRT with SRT did not show a statistical difference in terms of survival ( $p=0.530$ ), 3-y OS were 51.8%, 31.5% and 36.5%, respectively. In the univariate analysis, age  $<60$  years ( $p=0.901$ ), hormone receptor status ( $p=0.494$ ), number of brain metastasis ( $p=0.395$ ), extracranial metastasis ( $p=0.992$ ), recursive partitioning analysis (RPA) ( $p=0.394$ ), initial brain metastasis velocity (iBMV) ( $p=0.122$ ) did not affect OS. The median time to local treatment failure (LTF) was shorter in WBRT group, the median time to distant treatment failure (DTF) was shorter in SRT group.

**Conclusion:** This single-center series of consecutive patients with brain metastasis from breast cancer treated with SRT had a similar OS when compared to WBRT, consistent with the literature. However, large number of patients are needed to demonstrate important parameters affecting survival.

**Keywords:** Brain metastasis; breast cancer; stereotactic radiotherapy

	<b>SRT</b> (n=17)	<b>WBRT</b> (n=17)	<b>SRT+WBRT</b> (n=11)
<b>LTF</b>			
Patient (n)	10	11	4
Time, (median)	12.5 (6-26)	7 (4-17)	22.5 (4-36)
<b>DTF</b>			
Patient (n)	9	10	3
Time, (median)	10 (3-18)	12 (6-24)	24 (16-36)
<b>Reirradiation</b>			
Primary lesion	2	7	1
New lesion	4	6	1
Both	3	4	1

LTF: Local Treatment Failure; DTF: Distant Treatment Failure



# Epithelial-mesenchymal transition and Cancer

**Dr. Elif Önder**

*Pamukkale University, Medical Faculty, Denizli, Turkey  
E-mail: mrsocerebrum@hotmail.com*

## **Abstract**

EMT (Epithelial Mesenchymal Transition) is a process in which epithelial cells lose their apical-basal polarity and cell-cell adhesion and become spindle-shaped mesenchymal cells with higher migration ability. EMT plays a role in both physiological processes such as embryogenesis and wound healing and pathological processes such as fibrosis and cancer. Mesenchymal cell differentiation of epithelial cells is through key transcription factors that serve as major regulators of cell-cell adhesion, cell polarity and motility. Cancer cells are often referred to as partial or transient EMT, where various combinations of epithelial and mesenchymal markers coexist. EMT activation has been associated with the acquisition of stem cell properties and identified EMT as a critical regulator of cancer stem cells. EMT guides the tumor initiation and metastasis capacity of cancer cells as well as increasing resistance to chemotherapy and immunotherapy. This review discusses the various regulatory mechanisms of EMT and its pathological roles in cancer.

**Keywords:** *Epithelial-mesenchymal transition; Stemness; Metastasis; Chemoresistance*

# Ethical Aspects of Embryo Research

**Assoc. Prof. E. Elif Vatanoglu-Lutz**

*Yeditepe University Medical Faculty, History of Medicine and Ethics Department, İstanbul/Turkey*

*E-mail: elif.vatanoglu@yeditepe.edu.tr*

## Abstract

The discovery, isolation, and culturing of human embryonic stem cells can be described as one of the most important break throughs in biomedicine of the century. However, human embryonic stem cell research is ethically and politically controversial because it involves the destruction of human embryos. Embryo research has always been at the core of discussions according to medical ethics principles. General ethics codes such as Nürnberg Code, Helsinki Declaration, European Biomedicine Convention have a lot of articles regarding this issue but still many debates all around the world are going on.

In 2018, a Chinese scientist claimed the birth of the two girls whose C-C chemokine receptor type 5 (CCR5) gene(s) were deleted by CRISPR/Cas9 [clustered regularly interspaced short palindromic repeats (CRISPR)/CRISPR associated (Cas)] technology. The reason for this experimental procedure was the father being infected with human immunodeficiency virus (HIV). The national and international authorities heavily criticized the scientist in terms of scientific pitfalls of the technique used and ethics and showed condemnation against him. There is still an urgent need for specific legal regulations in international, national, and institutional levels and law enforcement to control human germline gene editing based on the currently available techniques such as CRISPR/Cas9.

**Keywords:** Human embryonic stem cell, embryo research, gene editing, ethical aspects

## From Tuhfetü'l Tıb\* to the Medicine of the Future

Eser Epözdemir<sup>1</sup>, Assoc. Prof. E. Elif Vatanoglu-Lutz<sup>2</sup> and Prof. Bahar Uslu<sup>3</sup>

*Yeditepe University Medical Faculty History of Medicine and Ethics Department, Istanbul, Turkey*<sup>2</sup>

*Quinnipiac University, Frank Netter MD School of Medicine, Connecticut, USA*<sup>3</sup>

*E-mail*<sup>1</sup>: eserepozdemir@gmail.com, *E-mail*<sup>2</sup>: elif.vatanoglu@yeditepe.edu.tr, *E-mail*<sup>3</sup>: bahar.uslu.md.phd@gmail.com

### Abstract

Eser Epözdemir is a researcher artist and during her attendance to the residency programme of Air Bayrampaşa in 2018, she made a research about Maltepe Military Hospital (Asâkir-i Mansûre: the Maltepe Military Hospital-1826) which was built by the order of Sultan Mahmut II. The architect of this hospital was the famous Baylan family who had big reputation in architecture those days. During her deeper research, Eser Epözdemir found about the publishing house in the hospital and a magazine called Tuhfetü'l Tıb was published. Tuhfetü'l Tıb is in Ottoman language, it means “About Medicine” in English and “Tıbba Dair” in Turkish. It was mainly about mother and baby health. Eser Epözdemir started to think what the content of this magazine would be if it was published today. Then, she had interactions with Dr. Elif Vatanoglu-Lutz who is the founder of Oksitosin Medicine and Art Platform. Together, they started to make exhibition talks about the important medical subjects for the future. Their main wish is seeing more culture and art activities in health establishments mainly for the resilience of healthcare professionals. In this paper, they are writing about the future topics of medicine when/if Tuhfetü'l Tıb is published in the future.

**Keywords:** *Tuhfetü'l Tıb, Asâkir-i Mansûre: the Maltepe Military Hospital, medical ethics, medical history, futuristic medicine.*

\* Tuhfetü'l Tıb is in Ottoman language, it means ‘About Medicine’ in English and ‘Tıbba Dair’ in Turkish.

# Medically Assisted Reproduction in Turkish Law

**Assoc. Prof. Emel Badur**

*Çankaya University Faculty of Law, Ankara, Turkey  
E-mail: badur@cankaya.edu.tr*

## Abstract

The right to reproduce is defined as the ability of individuals and couples to freely decide on the number of children and their birth interval, have the necessary information for this, and not be subjected to any oppression and discrimination during their access to healthcare services. Medically assisted reproduction is directly related to right to reproduce.

The primary regulation regarding medical assisted reproduction in Turkish Law is included in the Annex 1 article of the Code on Organ and Tissue Removal, Preservation, Vaccination and Transplantation. In this regulation it is as follows:

“In cases where children cannot be conceived by natural means or there is medical necessity, reproductive cells or embryo can be applied to the mother candidate by making the female and/or male reproductive cells suitable for fertilization by medical methods and insemination inside or outside the body. This method is performed only between married spouses. These treatment applications can be carried out only by physicians authorized by the Ministry and in medically assisted reproductive treatment application centers licensed by the Ministry, within the framework of the medical principles determined by the Ministry. The principles and procedures for opening, operating and supervising medically assisted reproductive treatment application centers are regulated by a regulation issued by the Ministry.

It is forbidden to have a child and to be a surrogate mother through the implementation of reproductive cells taken from one or both spouses and the embryo obtained from these cells to be used for other people.

Donation using someone else’s reproductive cell and/or embryo and donating, selling, keeping, using, storing, transporting, importing, exporting and agency these transactions are prohibited.”

In terms of Turkish Law, the detailed regulation on the subject has been made by secondary legislation and the “Regulation on Assisted Reproductive Treatment Practices and Assisted Reproductive Treatment Centers” (Regulation) has been issued.

Within the scope of this Code and Regulation, it is possible to consider medically assisted reproductive applications in narrow and wide scope. These narrow-scope applications can be defined as a set of medical interventions aimed at making married couples have children. Medically assisted reproduction in the broad sense includes the reproduction of couples as well as the removal and storage of individuals’ reproductive cells or tissues. In accordance with the regulation made by the Code, the authority to perform these medical interventions is only granted to physicians.

In the 14th article of Biomedicine Convention, which is binding in terms of Turkish Law, a restrictive regulation regarding medically assisted reproduction applications has been included as follows: “The use of techniques of medically assisted procreation shall not be allowed for the purpose of choosing a future child’s sex, except where serious hereditary sex-related disease is to be avoided.”

**Keywords:** *Right to reproduce, Medically assisted reproduction, Turkish Law.*

# The Cytotoxicity Effects of Metformin and Lithium Substances in the Human Intestinal Caco-2 Cells

**Emine Tural, MD<sup>1</sup>**

*Medeniyet University, Faculty of Medicine, Department of Histology and Embryology, Istanbul, Turkey<sup>1</sup>  
E-mail<sup>1</sup>: eminetural@gmail.com*

## Abstract

**Introduction:** Caco-2 is a human epithelial cell line frequently used as a model of the intestinal epithelial barrier. The Caco-2 cell line is derived from a colon carcinoma. Colorectal cancer is the third most common cancer worldwide. Many drugs are used clinically but still lack of effective therapy so far. The anticancer effects of metformin have been discussed and clinical studies are still ongoing in many areas such as endometrial cancer and colorectal cancer. Previous studies with lithium have shown that it has antineoplastic effects in various cancers including colorectal cancer, gastric cancer, and neuroblastoma.

**Aim:** The purpose of this study to is to determine of cytotoxicity effects of metformin and lithium substances in the human intestinal Caco-2 cells according to dose- dependent manner.

**Materials and methods:** Human intestinal Caco-2 cells were cultured in DMEM medium supplemented with 10% FBS, 20 units/ml penicillin and 20 µg/ml streptomycin and maintained in a humidified atmosphere of 95% air and 5% CO<sub>2</sub> at 37 °C. Cytotoxic activity of metformin and lithium in Caco-2 cells were determined by using XTT assay according to manufacturer's instruction. Cells were treated with different concentrations of metformin and lithium. Formazan formations were quantified spectrophotometrically at 450 nM (reference wavelength 630 nM) using a microplate reader. IC<sub>50</sub> values were determined by AAT Bioquest online IC<sub>50</sub> Calculator.

**Results:** The decrease in cell viability of the colorectal carcinoma Caco-2 cells was observed corresponding to increasing dose of metformin and lithium. In this study, cell viability and IC<sub>50</sub> values of metformin and lithium in Caco-2 cells were determined as respectively 26.1592 mM at 24th hour and 145.3614 µM at 48th hour.

**Conclusion:** These results indicated that metformin and lithium could be a potential therapeutic agent in colorectal carcinoma and further studies are required to determine probabilities.

**Keywords:** *Colorectal Carcinoma, Caco-2 cells, Metformin, Lithium*

# A Comparison of Propofol with Ketofol For Sedation Quality and Side Effects in Patients Undergoing Colonoscopy

Ender Çam, MD<sup>1</sup>, Prof. Deniz Karakaya<sup>2</sup> and Prof. Sibel Barış<sup>3</sup>

*Department of Anesthesiology and Reanimation, Gazi State Hospital, Samsun, TURKEY<sup>1</sup>*

*2Department of Anesthesiology and Reanimation, Ondokuz Mayıs University Faculty of Medicine, Samsun, TURKEY<sup>2,3</sup>*

*E-mail<sup>1</sup>: aesire@hotmail.com, E-mail<sup>2</sup>: dkarakaya65@gmail.com, E-mail<sup>3</sup>: sbaris@omu.edu.tr*

## Abstract

Colonoscopy is an endoscopic method and it is better to perform this procedure under sedoanalgesia in order to eliminate patients' anxiety, the colic-like pain and discomfort that occur during the procedure [1].

The aim of this study was to compare the effects of propofol and propofol+ketamine (ketofol) on sedation and side effects in patients undergoing colonoscopy.

**Material-Method:** 50 patients with ASA I-II that are between the ages of 18-65. The patients in the propofol group and Ketofol group were given 0.1mL/kg of drug and/or combination of drugs. The vital parameters, injection pain, spontaneous time of opening eyes, the time of Modified Aldrete Score (MAS)  $\geq 9$  and the amount of medication used during the procedure and in the recovery room were recorded.

**Results:** There was no significant difference between the two groups during the procedure and in the recovery room ( $p < 0.05$ ). It was shown that the ketamine reduces the amount of propofol by 50% and propofol induced injection pain. Ketofol had no positive effects on hemodynamic and respiratory parameters.

**Conclusion:** We assert that the ratio of combinations will vary depending on the necessary sedation level and analgesic need of the procedure to be performed and depending on the frequency of the administration of additional doses. Although ketofol is being used in different procedures and different age groups in the recent years, there is still need for studies conducted with different drug dosages of this combination.

**Keywords:** Colonoscopy, Ketofol, Hemodynamics, Sedoanalgesia

# International scientific collaborative activities: Barriers and opportunities

**Asst. Prof. Şule Başar<sup>1</sup> and Assoc. Prof. Enes Emre Başar<sup>2</sup>**

*Bayburt University, Faculty of Health Science, Nutrition and Dietetics Department, Bayburt, Turkey<sup>1</sup>*

*Anadolu University, Faculty of Business Administration, Eskişehir, Turkey<sup>2</sup>*

*E-mail<sup>1</sup>: sbasar@bayburt.edu.tr, E-mail<sup>2</sup>: eebasar@anadolu.edu.tr*

## Abstract

As known, Science is increasingly global and becoming a global enterprise. Cross-border flows of knowledge are increasingly common and frequent. The percentage of internationally co-authored science publications rose from 16.7% to 21.7% in the period 2010–2020. Current scientific research is entering an era of international collaboration, which is becoming the key force in leading-edge scientific discovery and producing a large number of high-quality research results. At the same time, the rise of the research network is changing the global balance of science and reshaping the global scientific landscape.

In the age of the globalizing knowledge economy, international knowledge flows have increasingly become a key driving factor for economic growth and competitiveness. Both advanced and developing countries will benefit from the globalization of science, which is a ‘win-win’ game. In addition, globalization will bring about specialization and networking of science activities, increase the interconnectedness and interdependence between countries, and result in the emergence of an international collaboration network. The globalization of science may be explained by the following reasons. First, the globalization of science is similar to economic globalization, which is a result of the international division of labor among researchers, and scientific activity is increasingly coordinated at the global scale. Second, new scientific discovery or innovation is not a simple linear input-output process: the nature is complex and nonlinear because it is full of uncertainty and risk. Third, with the increased specialization of knowledge production, closed innovation has shifted to an open innovation paradigm. Fourth, humanity faces many overwhelming global challenges, such as Covid 19 pandemic, climate change, energy security, pollution and so on, which are interdependent and interrelated. Fifth, rapid progress in transportation and Information Communication Technology, such as high-speed Internet, has reduced the cost of travel and communication, which makes cross-border collaboration a possibility.

The globalization of science is manifested in various forms, such as international co-patenting networks, international patent citation networks, international co-authorship networks, international publication citation networks, international academic conferences, global talent mobility networks and R&D collaboration networks. The geography of international scientific collaboration is highly concentrated and the whole network is dominated by certain European nations, China and the USA. The rise of emerging scientific nations is breaking the global balance of science. The geography of global science has evolved from a bipolar world to a multi-polar world. At the same time, the mean center of gravity of global knowledge production is shifting from the West towards the East and from the North to the South.

Over the past two decades, R&D funding has increased globally, contributing to tremendous growth in global science publication output. The pursuit of R&D can be framed as a competition between different nations, which at times has been a driving force in promoting policy change and encouraging increased national expenditures in science and technology R&D. However, many scientists are approaching science with an increasingly collaborative outlook. While the United States

still maintains its position as the top R&D contributor, China has steadily increased expenditures and publication outputs leading the world in total publication since 2016. This increase in research has built a worldwide network of scientists participating in intra-national and international collaboration.

**Keywords:** *The globalization of science; scientific collaboration; globalizing knowledge economy*



# Clinical outcomes of intraarticular PRP and corticosteroid combination in advanced osteoarthritis

**Erhan Okay, MD**

*Istanbul Goztepe Prof. Dr. Suleyman Yalcin City Hospital, Department of Orthopaedics and Traumatology  
E-mail: erhanokay@yahoo.com*

## Abstract

**Background:** Advanced osteoarthritis is affecting the elderly population, which reacts via catabolic and inflammatory joint environment. There is limited data about intraarticular combination of PRP and steroids on clinical outcomes of advanced osteoarthritis.

**Aim:** The objective of this study is to compare the clinical response of corticosteroid (CS) and platelet-rich plasma (PRP) treatment in 2 groups of patients affected by advanced osteoarthritis

**Methodology:** A total of 68 patients affected by clinically and radiographically documented with grade 4 gonarthrosis according to the Kellgren-Lawrence classification were included in this study. The patients were identified into 2 study groups. Thirty-two patients (Group 1) received 3 intra-articular injections of PRP (5mL) and steroid (1 cc – 5 mgr triamcinolone). Thirty-six patients (Group 2) received 3 intra-articular injections of PRP with one-week interval (5 mL). An unblinded physician performed injection once a week for 3 weeks into the affected knee in both groups. All patients were evaluated with the VAS score and the Knee injury and Osteoarthritis Outcome Score (KOOS) subscales before the infiltration, at 2 and 6 months after the first injection.

**Results:** Two groups are similar with regard to demographic variables (age, gender, BMI, stage of arthritis). The combination of intra-articular PRP with steroids resulted in a significantly superior clinical outcome, with sustained lower VAS ( $p < 0.01$ ) and improved KOOS subscales ( $p < 0.01$ ) except for KOOS sporting activity and quality of life within 6 months compared to intra-articular PRP only injection.

**Conclusion:** Treatment with intra-articular PRP and steroids showed a significantly better clinical outcome than did treatment with PRP, with sustained better KOOS scores

**Keywords:** *Platelet rich plasma; steroid; intraarticular injection; advanced osteoarthritis*

# A COVID-19 case complicated by ecchymosis and cyanosis

**Dr. Esmeray Mutlu Yılmaz<sup>1</sup>**

*Clinic of Clinical Microbiology and Infectious Diseases<sup>1</sup>  
Health Sciences University, Samsun Training and Research Hospital, Samsun, Turkey<sup>1</sup>  
E-mail<sup>1</sup>: emutlu55@gmail.com*

## **Abstract**

It has been emphasized that the main causes of death of COVID-19 patients are DAD (Diffuse Alveolar Damage) and DIC (Disseminated intravascular coagulation). Microthrombosis is an important clinical feature of COVID-19 and was seen in 91.3% of patients who died. Here, a case of COVID-19 complicated by ecchymosis and cyanotic appearance due to microthrombus developing in superficial veins is presented.

A 57-year-old female patient presented with bruising and pain in her feet and right breast. She had COVID-19 PCR positivity 10 days before and used the favipiravir treatment at home for five days. Two days after the treatment ended, she complained of sudden onset and increasing pain in her right breast and feet. She applied to the emergency department because of the increasing bruising in the painful areas. At the time of admission, her general condition was moderate to poor, saturation was between 85-90% at room air. Significant coldness and bruising were observed in the entire right breast, back and both feet. In thorax CT examination; widespread signs of infiltration were observed in both lungs consistent with viral pneumonia. In bilateral lower extremity arterial doppler examination; no pathology was detected. In venous doppler examination; the right saphenous parva was observed in the subacute period with thrombus occlusion. Her urea and creatine were elevated. Dobutamine, perlinganit and M infusion was started. Prednol 120mg/d IV was initiated in consultation with rheumatology. In peripheral smear; 90% neutrophils, toxic granulation in the myeloid series and reactive thrombocytosis with a left shift were observed. Thereupon, ceftriaxone 2x1 g IV was started empirically. Enoxaparin sodium 2x0.6 sc and acetylsalicylic acid 100 mg/d were started.

COVID-19 thrombosis includes macro and microthrombosis due to coagulation and fibrinolysis markers. In this case, ecchymotic lesions on the breast and back were thought to be due to microthrombi associated with COVID-19. Antiagregant use and thrombocytopenia should be considered in the differential diagnosis. Our patient did not use anticoagulants and her thrombocyte counts were normal. Acute Kidney Injury is a common complication of severe COVID-19 infection. Ischemic acral lesions have been reported in patients with COVID-19 in the literature. It has been stated that these lesions are thrombotic lesions that occur as a result of treatment or COVID-19. When skin lesions, lung and kidney involvement developed in this patient are evaluated together; hypotension and hypoperfusion cannot be explained by classical sepsis, diffuse intravascular coagulation, vasculitis, endocarditis, or severe peripheral arteriosclerosis. Clinical findings support microthrombosis interpreted as activation of the coagulation system associated with COVID-19.

**Keywords:** COVID-19, microthrombosis, ecchymosis, cyanosis

# The frequency of micronodular type neuroendocrine cell hyperplasia detected in gastric endoscopic biopsies and its relationship with clinicopathological parameters

Oğuzhan Okcu<sup>1</sup>, Ezgi Hacıhasanoğlu<sup>2</sup> and Bayram Şen<sup>3</sup>

*Recep Tayyip Erdogan University, Medical Faculty, Department of Pathology, Rize, Turkey<sup>1</sup>*

*Yeditepe University, Medical Faculty, Department of Pathology, Istanbul, Turkey<sup>2</sup>*

*Recep Tayyip Erdogan University, Medical Faculty, Department of Biochemistry, Rize, Turkey<sup>3</sup>*

*E-mail<sup>1</sup>: oguzhanokcu@hotmail.com, E-mail<sup>2</sup>: ezgihaci@yahoo.com, E-mail<sup>3</sup>: byrmnsn06@gmail.com*

## Abstract

Neuroendocrine cell hyperplasia (NECH) is defined as linear or micronodular clusters of at least 5 neuroendocrine cells. By definition, micronodular clusters should be  $\leq 150$  microns in greatest dimension. In our study, we aimed to determine the rate of micronodular type NECH in gastric endoscopic biopsy cases and to determine the pathological and clinical features that may be associated with this condition.

Among 3974 gastric endoscopic biopsy materials examined between July 2016 and January 2021, 38 patients diagnosed with micronodular type NECH were included in the study. Thirty-nine patients without micronodular type NECH were used as a control group. Clinical information (age, gender and history of proton pump inhibitor use) of the cases was obtained from the hospital electronic database. The presence of inflammation, activation, glandular atrophy, intestinal metaplasia, and *Helicobacter pylori* evaluated in routine pathological examination, as well as clinical features; age, gender, history of proton pump inhibitor (PPI) use were compared among two groups by using statistical analysis (Chi-square test, Pearson Chi-Square, Fisher's Exact Test, Spearman correlation analysis).

In our study, micronodular type NECH was found in 38 cases (0.95%) in gastric endoscopic biopsy samples belonging to 3974 different patients. Twenty-two patients were female, 16 were male. The age range was 33-88 and the average age was 62. Thirty-one out of 38 cases had one or more use of PPI. PPI use was not detected in 7 patients. Among cases without NECH, 17 of the cases were female, 22 of them were male. The age range was 18-90 and the average age was 53. In 23 of 39 cases, there was one or more history of PPI use, and no PPI use was found in 16 patients. In statistical analysis, the use of PPI was found with a significantly higher rate in patients with micronodular type NECH. The presence of intestinal metaplasia and glandular atrophy showed borderline statistical significance between groups. There was no significant difference in age and gender distribution between the groups.

Micronodular type NECH is rarely encountered in gastric endoscopic biopsy materials, the main reason of which is being reported as the use of proton pumps. Supportively, in our study, a statistically significant relationship was observed between the use of PPI and the development of micronodular type NECH. If the history of PPI use could be found out in all gastric endoscopic biopsies and neuroendocrine markers could be applied in all cases, a more definite conclusion could be reached about the relationship between the development of NECH and PPI use. In order to fully understand the clinical significance and reason of this finding, studies involving much larger numbers of cases and with long-term follow-up are required.

**Keywords:** *Micronodular neuroendocrine cell hyperplasia; proton pump inhibitor; gastric biopsy*

## Renal Tissue Morphology and Morphometric Alterations with Different Fixatives (Formalin, Bouin's, Alfac and B5 Solutions)

**Dr. Fatma Mert<sup>1</sup>, Dr. Bilge Serdaroğlu<sup>2</sup>, Dr. İbrahim Alptekin<sup>3</sup>,  
Asst. Prof. Ferda Topal Çelikkan<sup>4</sup> and Prof. Oya Evirgen<sup>5</sup>**

Ankara University, Medical Faculty, Department of Histology and Embryology, Ankara, Turkey <sup>1,2,3,4,5</sup>  
E-mail <sup>1</sup>: f.mert417@gmail.com, E-mail <sup>2</sup>: blgsrdrgl@gmail.com, E-mail <sup>3</sup>: ibrahimalptekin4221@gmail.com,  
E-mail <sup>4</sup>: ferdatopal@gmail.com, E-mail <sup>5</sup>: oya.evirgen@gmail.com

### Abstract

**Introduction:** Fixation is a critical step to preserve the tissue morphology and cellular details for histologic examination. The fixatives have advantages and disadvantages including the tissue shrinkage and the quality of histochemical staining. Formalin is mostly used fixative for study and research in the field of anatomy, histology and pathology [1-5]. In the present study, we aimed to evaluate the diameter of renal corpuscle, overall staining characteristics and cellular details of the renal tissue fixed with formalin, Bouin's, alfac and B5 solutions under the light microscope.

**Methods:** This study was conducted in Ankara University, Faculty of Medicine Department of Histology and Embryology. Twelve (n = 12) adult BALB/c mice were randomly assigned to 4 groups. The animals were sacrificed and bilaterally the kidneys were fixed with 4% formalin, Bouin's, alfac and B5 solutions. After fixation kidney tissues were dehydrated in graded series of alcohol and embedded in paraffin. 4 µm paraffin sections were stained with hematoxylin and eosin, periodic acid Schiff and Mallory Azan. For morphologic analysis the kidney tissue sections of all animals in each fixative group were evaluated according to four parameters: cytoplasmic detail, nuclear detail, tissue integrity and overall staining were scored as (1: poor, 2: moderate, 3: good, 4: excellent). For morphometric analysis in each fixative group digital 8-bit RGB images were captured by a AxioCamMRC mounted on the light microscope under a 40x magnification lens. A total of 20 randomly selected renal corpuscle in cortex region from each animal, the digital images were assessed for renal corpuscle diameter in a blinded by two senior histologist using ImageJ program (<http://rsb.info.nih.gov/ij/>).

**Results:** All of the fixatives that were used demonstrated that kidney tissues were well preserved. The cortex and medulla regions were easily distinguished and the integrity of these regions were maintained. The cell nuclear and cytoplasmic details of the nephron and its parts were clearly visible. The cellular details (cytoplasmic details, nuclear details) and tissue integrity that were evaluated and scored revealed that alfac and Bouin's solutions gave more satisfactory results on the mice kidney tissue morphology than formalin and B5 solutions. The overall histologic score showed that Bouin's, alfac and formalin fixatives were better than B5 fixative. The morphometric analysis demonstrated that the renal corpuscle diameter was moderately increased in B5 group ( $88.59 \pm 12.5$  µm;  $p < 0.05$ ) compared with the other fixative groups Bouin's, alfac, formalin ( $80.58 \pm 12.38$  vs  $82.18 \pm 9.75$  vs  $81.86 \pm 9.65$ ) and the difference was statistically significant.

**Conclusion:** In conclusion, the kidney tissue morphometric and morphologic analyses suggested that the fixatives like alfac and Bouin's which contains glacial acetic acid were more effective in preserving kidney tissue architecture than formalin and B5 fixation solutions.

**Keywords:** Fixation, Fixatives, Renal tissue, morphology, morphometry.

## New challenge in adolescents: ‘at risk polycystic ovary syndrome’ estimation

**Dr. Fatma Nurgül Taşgöz<sup>1</sup>**

*Health Sciences University, Bursa Yüksek İhtisas Training and Research Hospital, Bursa, Turkey<sup>1</sup>  
E-mail<sup>1</sup>: fna78ahoo.com*

### Abstract

**Background:** Polycystic ovary syndrome (PCOS) is a common endocrine disorder seen 8-13% in reproductive age and 6-18% in adolescence. Although the Rotterdam criteria; when 2 of the 3 diagnostic criteria are present. 1) signs of clinical or biochemical hyperandrogenism (2) oligo/anovulation and/or (3) polycystic ovaries are diagnostic for women of reproductive age, the diagnosis of PCOS in adolescents is challenging. Because clinical features can overlap pubertal developmental events. Due to the immaturity of the hypothalamic-pituitary-ovarian axis in adolescents, menstrual irregularities with varying cycle length, comedonal acne and large multicystic ovaries are a common finding. Although there are no defined criteria for the diagnosis of PCOS in adolescent, Anti-Mullerian Hormone (AMH) levels and pelvic ultrasound for polycystic ovarian morphology (PCOM) not recommended for PCOS diagnosis in current guidelines. PCOS diagnosis should be made in the presence of both persistent oligomenorrhea and clinical or biochemical hyperandrogenism.


**Purpose:** To address adolescent girls at risk of PCOS; presenting with isolated symptoms such as irregular menstrual cycles or clinical hyperandrogenism, for which the diagnosis of PCOS is uncertain. Beside this to compare the clinical and laboratory characteristics of at risk of PCOS adolescents with their healthy control and with definite PCOS diagnosed peers.

**Method:** 98 adolescent girls who admitted to our gynecology outpatient clinic those had menstruation cycles for at least 3 years or more were recruited. Adolescents with other disease causing androgen excess and ovulatory dysfunction, those treated for PCOS were excluded.

Menstrual intervals shorter than 20 days or longer than 45 days were oligoanovulation. Moderate to severe inflammatory acne, seborrhea, modified Ferriman Gallaway score above 8 were clinical hyperandrogenism. AMH, 17-hydroxyprogesterone, LH, Total testosterone, free testosterone, TSH, PRL, DHEAS levels were measured, BMI, HOMA-IR and LH/AMH were calculated. All participants were evaluated by pelvic ultrasound.

**Results:** 59 of 98 adolescents were diagnosed with PCOS according to Rotterdam Criteria. When the same 98 girls were evaluated according to current guidelines 37 of them were diagnosed with PCOS, 40 girls were in the at risk PCOS group, and 21 girls who did not have any of the 3 criteria included in the control group (Figure 1). There were no differences between the groups in terms of age, BMI, 17 hydroxyprogesterone, LH, Total testosterone, free testosterone, TSH, PRL, DHEAS and HOMA-IR. AMH levels were significantly higher ( $p < 0,001$ ) and LH/AMH ratios were significantly lower ( $p < 0,001$ ) in PCOS and at risk of PCOS group than control group (Table 1).

**Conclusion:** While avoiding overdiagnosis of PCOS in adolescents, there is also a possibility of underdiagnosis. AMH values may be high even without polycystic ovarian morphology. AMH levels and LH / AMH ratio may be useful in guiding patients at risk of PCOS to clinical follow-up.



**Figure 1.** Distribution of cases according to clinical and diagnostic features

**Table 1.** Clinical characteristics of the groups divided according to the diagnosis of PCOS in adolescents in the light of current guidelines.

	Group 1 PCOS (n= 37)	Group 2 at risk PCOS (n= 40)	Group 3 Control (n= 21)	p
Age(years)	18 (15-19)	18 (15-19)	18 (16-19)	0,574
BMI	24,7 (18,2- 41,5)	24,9 (18- 36,8)	24,8 (19,3- 38,3)	0,782
17 OH Progesteron	1(0,01- 2,92)	1,075 (0,01- 3,10)	1(0,33- 2,52)	0,223
AMH	9,13(0,11- 26,9) <sup>a</sup>	6,38(0,01- 27,80) <sup>a</sup>	2,32(0,01- 10,60)	<0,001
LH	5,19(0,10- 49,10)	5,94 (0,69- 19,51)	5,11 (2,22- 31,36)	0,977
Total testosterone	48(11-171)	39,61 (5,26- 210)	39 (8-68)	0,094
Free testosterone	2,53 (0,46-14,10)	2,25(0,18-14,49)	1,72 (0,54- 10,37)	0,365
Prolaktin	15,19 (2,85- 63)	16,70 (5,59- 68,10)	16,54 (2,24- 34,99)	0,954
TSH	2,12 (0,52- 4,70)	1,99 (0,39- 8,71)	2,08 (1,03- 3,90)	0,879
DHEAS	280,7 (19- 521)	296,6 (28,7- 744,2)	225,8(111,9- 311,6)	0,067
HOMA-IR	2,45 (0,65- 8,70)	2,25(0,94- 7,38)	2,09 (0,81- 5,88)	0,433
LH/AMH	0,72 (0,01- 15,11) <sup>a</sup>	0,79 (0,03- 9,16) <sup>a</sup>	2,05 (0,61- 14,34)	<0,001

Data presented as Median (min-max). BMI:body mass index; AMH:anti- mullerian hormone LH; *luteinizing hormone* TSH; thyroid stimulating hormone DHEAS: *dehydroepiandrosterone* sulphate HOMA-IR: homeostasis model assessment of insulin resistance

<sup>a</sup>There was a significant difference with compared group 3 in pairwise comparisons.

\*Independent Samples Kruskal- Wallis test was used. A p value of 0.05 was considered significant.

# Determination of Cell Viability in LNCaP Prostate Cancer Cells Treated with Boric Acid

Gizem Kabasakal, MD<sup>1</sup> and Mücahit Seçme, PhD<sup>2</sup>

*Pamukkale University, Faculty of Medicine, Department of Histology and Embryology, Denizli, Turkey<sup>1</sup>*

*Pamukkale University, Faculty of Medicine, Department of Medical Biology, Denizli, Turkey<sup>2</sup>*

*E-mail<sup>1</sup>: gizemkabasakal7@gmail.com, E-mail<sup>2</sup>: mehtersecme@gmail.com*

## Abstract

**Introduction:** Prostate cancer is the most diagnosed malignancy in men. Common methods of prostate cancer treatment include chemotherapy, hormonal therapy or gene therapy. Boric acid (H<sub>3</sub>BO<sub>3</sub>) is a weak acid, the physiological pH form of Boron. It is often used as an antiseptic, antifungal and antiviral.

**Aim:** The aim of the study was to demonstrate the cytotoxicity effects of boric acid in LNCaP prostate cancer cell lines in a dose-dependent manner.

**Materials and methods:** LNCaP prostate cancer cells were cultured in DMEM medium supplemented with 10% FBS, 20 units/ml penicillin and 20 µg/ml streptomycin and maintained in a humidified atmosphere of 95% air and 5% CO<sub>2</sub> at 37 °C. Cytotoxic and anti-proliferative activity of boric acid in LNCaP cells were determined by using XTT assay according to manufacturers' instruction. Cells were treated with different concentrations of boric acid. Formazan formations were quantified spectrophotometrically at 450 nM (reference wavelength 630 nM) using a microplate reader. IC<sub>50</sub> values were determined by AAT Bioquest online IC<sub>50</sub> Calculator.

**Results:** Decrease in cell viability of the LNCaP prostate cancer series was observed in direct proportion to the increase in boric acid dose. When cell viability was evaluated in LNCaP cells, the IC<sub>50</sub> value of boric acid was determined as 89.6713 µM at the 48th hour. These findings showed that boric acid could be a potential therapeutic agent for further studies.

**Keywords:** *Boric acid, LNCaP, Prostate Cancer*

# Comparison of Insulin Treatment with Mesenchymal Stem Cell Treatment in Experimental Type 1 Diabetes-Induced Rats

Gökçen Gökçe<sup>1</sup>, Murat Tosun<sup>2</sup>, Hasan Hüseyin Demirel<sup>3</sup> and Esra Aslan<sup>4</sup>

Ankara University, Faculty of Medicine, Histology-Embryology Department, Ankara/ Turkey<sup>1</sup>

Afyon University of Health Sciences, Faculty of Medicine, Histology-Embryology Department, Afyon/ Turkey<sup>2</sup>

Afyon Kocatepe University, Faculty of Veterinary Medicine, Pathology Department, Afyon/ Turkey<sup>3</sup>

Afyon University of Health Sciences, Faculty of Medicine, Histology-Embryology Department, Afyon/ Turkey<sup>4</sup>

E-mail<sup>1</sup>: gkgkcn@hotmail.com, E-mail<sup>2</sup>: drmtosun@yahoo.com, E-mail<sup>3</sup>: hdemirel@aku.edu.tr, E-mail<sup>4</sup>: dr\_esragul@hotmail.com

## Abstract

**Introduction:** Diabetes is one of the leading causes of mortality and morbidity worldwide, with damage to beta cells. Mesenchymal stem cell (MSC) treatment is one of the alternative methods instead of insulin. The aim of this study is to investigate the effects of MSCs on Streptozotocin-induced diabetes in male albino rats, their potential therapeutic effect and possible applications in human diseases.

**Method:** In this study, Wistar albino rats were used. Five groups were formed. The diabetic model was created by intraperitoneal injection of Streptozotocin (STZ). MSC was injected intrapancreatically in determined groups, and all experimental groups were sacrificed three weeks later. During the experiment, blood glucose and insulin levels of all subjects were measured regularly.

**Results:** In the diabetic rats, it was seen that there was no significant change in the structure of the stroma in the group given MSC alone and the new vascular formations were more pronounced than the group with only STZ and insulin applied. In the examination of pancreatic tissues of diabetic rats treated with MSC and insulin, the stroma was homogeneous, neovascularization was increased, the structure and cellular structure of Langerhans islets were better than in other groups. In the examination, it was observed that there was no necrotic degenerative acinar cells or islet cells and edema decreased in the acinus structure and neovascularization became evident.

**Conclusion:** In addition to insulin therapy in DM treatment, it has been observed that intrapancreatic MSC therapy can be used as an additional treatment for gaining healthy pancreatic morphology.

**Keywords:** *Mesenchymal Stem Cells, Streptozotocin, Insulin, Diabetic Rat*



# Oxidative Effects and Biochemical Markers Related to Disease Severity Parameters in Cystic Fibrosis Cases

**Dr. Göksenin Ünlügüzel Üstün**

*Health Sciences University, Samsun Training and Research Hospital, Samsun, Turkey  
E-mail: mdgoksenin@hotmail.com*

## Abstract

Cystic fibrosis (CF) is the most common lethal autosomal recessive disorder seen in Caucasians. The major clinical manifestations of CF are exocrine pancreatic deficiency, male infertility, and chronic respiratory diseases. Malnutrition, initiated by exocrine pancreatic deficiency and contributed to chronic infections, can lead to a decline in respiratory functions, increased risk of lung infections, and complications like decreased bone mineral density and latent puberty.

This study aimed to examine the relations between major scoring systems and respiratory function test results with biochemical markers and then identify the most suitable biochemical marker(s) for patient follow-up and show the relation between growth development and disease severity. Furthermore, we examined oxidative conditions caused by CF systemic component and their effects on disease severity and growth development.

Children who were followed up with a CF and non-CF bronchiectasis diagnosis in Marmara University Faculty of Medicine, Chest Diseases division of Pediatrics Department were included in the study after obtaining parental written consent. Then blood and sputum samples were collected from the children following 8 hours of fasting.

Thirty-one patients with CF ( $10,5 \pm 2,9$  yrs) and 16 non-cystic fibrosis bronchiectasis patients ( $10,5 \pm 2,3$  yrs) taken as the control group were enrolled in the study. There was no significant difference between CF and non-CF bronchiectasis group in lung status, determined by respiratory function test (Forced expiratory volume-FEV1) ( $p > 0.05$ ). Anthropometric measurements and bone mineralization density did not significantly differ between CF and control group ( $p > 0.05$ ). Biochemical inflammation marker C-reactive protein (CRP), sputum IL-8, TNF- $\alpha$ , monocyte respiratory burst results, and biochemical marker of growth-development IGF-1 and IGFBP-3 also showed no significant difference between CF and control group ( $p > 0.05$ ). Serum Malondialdehit (MDA) concentrations, showing systemic oxidative status, were significantly different between CF [ $1,74$  ng/mL ( $1,5-1,95$ )] and control group [ $1,34$  ng/mL ( $1,05-1,79$ )] ( $p < 0.05$ ). There were strong correlations between Shwachman, CT scoring systems, and FEV1 with CRP and albumin. Moreover, we demonstrated significant correlations between IGF-1 with Shwachman, CT scores, and FEV1, FVC. Similarly, IGFBP-3 significantly correlated with FEV1, FVC, free % fat, and free fat mass. Furthermore, there were correlations between monocyte oxidative burst with Shwachman, CT scores, and FEV1 ( $p < 0.05$ ). Likewise, a marker of systemic oxidative status, MDA significantly correlated with radius z score and weight. All these findings showed us systemic component of CF plays an important role in disease progression independently of lung status. We suggest that serum MDA could be used in the follow-up of oxidant/antioxidant status, resulting from systemic oxidative stress seen in CF.

IGF-1 and IGBP-3 levels seemed promising in deciding the time of GH supplement therapy. Thus, without affecting the initiation of cystic fibrosis-related diabetes, improvement in growth development and respiratory functions could be provided.

In conclusion, Shwachman and CT scoring systems seemed to be more efficient in determining disease severity than the other evaluated scoring systems. As all these scoring systems are impractical

and have low sensitivity, especially for children with CF, we recommended CRP usage with albumin in determining disease severity and monitoring treatment response.

*Keywords: Cystic fibrosis, biochemical markers, Disease Severity Parameters*

# Determination of Sleep Quality in Patients with Rotator Cuff Tears

Gonca Sağlam, MD

*Karadeniz Technical University, Medical Faculty,  
Department of Physical Medicine and Rehabilitation, Trabzon, Turkey  
E-mail: goncasaglam@hotmail.com*

## Abstract

**Background:** Sleep disturbances due to shoulder pain are quite frequently encountered complaints in daily clinical practice and impact physical, social, and emotional functioning of patient's quality of life, causing mood disorders and anxiety.

**Purpose:** This study aimed to assess the sleep quality, pain and functionality in patients who have been diagnosed with rotator cuff tear (RCT), and to compare RCT patients with healthy subjects in terms of sleep quality.

**Methodology:** Fifty-two patients with a diagnosis of RCT as determined by magnetic resonance imaging and fifty age and gender matched healthy individuals were enrolled in this cross-sectional study. Exclusion criteria included: shoulder surgery in the past 3 months, severe glenohumeral osteoarthritis and adhesive capsulitis, fracture history involving the shoulder region or glenohumeral joint dislocation, a documented history of sleep disorder, previously diagnosed mood or psychotic disorder. All participants completed a questionnaire regarding their demographic data and sleep quality evaluated by Pittsburgh Sleep Quality Index (PSQI). RCT patients were also examined using the Visual Analog Scale (VAS) and the American Shoulder and Elbow Surgeons scale (ASES). Statistical analyses were performed using the IBM SPSS Statistics 22 (IBM, SPSS, Turkey) software. Data represented as mean  $\pm$  standard deviation. Student's t test and the Mann-Whitney U test were used to compare quantitative data of the RCT patients with those of the control group. Associations between the VAS, ASES, and global PSQI scores were explored with Pearson correlation coefficient test. A p value of  $<0.05$  was considered statistically significant.

**Results:** Among all participants with RCT, the mean age was  $52.4 \pm 13.4$ , whereas the mean VAS score was  $6.4 \pm 2.5$  and the mean ASES score was  $48 \pm 24$ . The presence of nocturnal pain was determined in 92% of RCT patients. There was a significant difference between the RCT group and control group in terms of global PSQI scores and all subdivisions ( $p < 0.01$ ). The mean global PSQI score for the RCT group and the control group were  $11.08 \pm 5.14$  and  $4.70 \pm 2.66$ , respectively. The comparison of global PSQI scores, VAS and ASES scores revealed a significant correlation in the RCT group ( $r = 0.52/p < 0.01$ ,  $r = 0.54/p < 0.01$ , respectively).

**Conclusion:** This study highlighted that patients with RCT suffer from sleep disturbances as represented by poorer sleep quality index scores. Moreover, sleep quality was found to be correlated with pain and functionality. Patients with shoulder pain due to RCT may benefit from pain management and cognitive-behavioral interventions that specifically target sleep disturbances.

**Keywords:** *Pittsburgh Sleep Quality Index; Rotator cuff tears; Sleep quality.*

# Determination of the functions of miRNAs associated with PI3K-Akt signaling pathway and p53 signaling pathway in prostate cancer

Dr. Gözde Öztan<sup>1</sup>

*Istanbul University, Istanbul Faculty of Medicine, Department of Medical Biology, Istanbul, Turkey<sup>1</sup>  
E-mail<sup>1</sup>: gozdeoztan@istanbul.edu.tr*

## Abstract

Prostate cancer (PCa) constitutes a major health problem in Western countries. In our study, it was aimed to reveal the connections between miRNA and pathway associated with prostate cancer using miRwayDB database. In the study, the biological roles and related mechanisms of hsa-miRs in PCa have been demonstrated. When miRwayDB database results are evaluated; It has been determined that hsa-miR-124, which is located in the PI3K-Akt signaling pathway, disrupts the adhesion, migration and invasion of prostate cancer cells by down-regulation of TALIN 1. TALIN 1 was indicated as a potential target of hsa-miR-124. The miRNA profiling analysis results show that miR-188-5p is significantly downregulated in metastatic PCa. It has been found that overexpression of LAPTM4B is associated with PCa progression. hsa-miR-302a suppresses AKT expression which causes subsequent change in the PI3K-Akt signaling pathway in prostate cancer. hsa-miR-4638 suppresses castration resistance of prostate cancer via targeting KIDINS220 and PI3K-Akt signaling pathway. Loss of miR-4638-5p may lead to castration resistant PCa through the activity of KIDINS220 and PI3K/AKT pathway. hsa-miR-7 suppresses the stemness of prostate cancer stem-like cells and tumorigenesis through inhibiting KLF4/PI3K/Akt/p21 pathway.

Gene-gene and network interactions were created between 6 genes using the STRING analysis. A significant enrichment was detected in a subgroup of genes involved in the response to fluid shear stress [GO:0034405], negative regulation of protein kinase B signaling [GO:0051898], positive regulation of nitric oxide biosynthetic process [GO:0045429], cellular response to nerve growth factor stimulus [GO:1990090] in the biological process by Gene Ontology (GO) enrichment analysis. The linkage between the PI3K-Akt signaling pathway and p53 signaling pathway located on the KEGG pathway in prostate cancer was determined by means of the KLF4, AKT1 and MDM2 genes. Then, the characteristics of these three genes were compared using the Genecards Human Gene Database. According to this, KLF4 protein is thought to control the G1-to-S transition of the cell cycle following DNA damage by mediating the tumor suppressor gene p53. AKT1 (protein kinase B) is a known oncogene. Expressed in PCa and levels increase from the normal to the malignant state. MDM2 protein can promote tumor formation by targeting tumor suppressor proteins, such as p53, for proteasomal degradation. This gene is itself transcriptionally regulated by p53.

As a result, the elucidation of molecular pathways regulated by tumor suppressor miRNAs can shed light on oncogenic and metastatic processes in PCa. hsa-miR-7 has been identified as a new tumor suppressor miRNA and it abolishes the stemness of PCSCs and inhibits prostate tumorigenesis by suppressing an important stemness factor KLF4. AKT has been identified as a target gene in which hsa-miR-302a exerts its inhibitory role in PCa. hsa-miR-340 acts as a tumor suppressor via the p53 signaling pathway by inhibiting MDM2 expression in prostate cancer.

**Keywords:** Prostate cancer; miRNA; pathway

# Efficacy of Lacosamide Therapy in Focal Onset Refractory Epilepsy of Childhood: A Single Center Experience

**Dr. Halil Ural Aksoy<sup>1</sup>, Dr. Celil Yılmaz<sup>2</sup>, Dr. Senem Ayca<sup>3</sup> and Dr. Muzaffer Polat<sup>4</sup>**

*Sivas Numune Public Hospital, Department of Pediatric Neurology, Sivas, Turkey<sup>1</sup>*

*Celal Bayar University, Medical Faculty, Department of Pediatric Neurology, Manisa, Turkey<sup>2,4</sup>*

*Haseki Research Hospital, Department of Pediatric Neurology, Istanbul, Turkey<sup>3</sup>*

*E-mail<sup>1</sup>: uralaksoy@hotmail.com, E-mail<sup>2</sup>: clyilmaz@yahoo.com, E-mail<sup>3</sup>: Senemkaleci@yahoo.com,*

*E-mail<sup>4</sup>: polatmuzaffer@yahoo.com*

## Abstract

**Introduction:** Treatment of childhood refractory epilepsy is a challenge for clinicians. Lacosamide is a new generation antiepileptic drug which is being used for focal onset seizures of adults and children. Efficacy and safety of the drug for adults have been demonstrated in various studies. The aim of this retrospective cross-sectional study is to evaluate the efficacy and safety of lacosamide in childhood refractory focal seizures in our clinic.

**Methods:** We examined the medical records of 14 patients treated with lacosamide in our clinic between January 2016 and January 2020 in terms of demographic, etiological, neuroimaging findings, responses to treatment, adverse effects and drug-drug interactions. We evaluated the patients as responders to treatment whose seizure frequency decreased  $\geq 50\%$  after 6 months of lacosamide treatment.

**Results:** In 12 patients (85.7%) seizure frequency decreased  $\geq 50\%$  ( $p < 0.001$ ) while 5 of them (35.7%) were seizure free. Despite the long-term treatment one patient did not respond to lacosamide treatment, and 1 patient's treatment stopped due to increased seizure aggravation after lacosamide initiation. Clinical adverse effects were observed in 3 (21.4%) patients. Cardiac adverse effects or drug-drug interactions were not observed in any patient.

**Discussion and Conclusion:** We think that lacosamide is an effective and reliable treatment option for refractory focal seizures of childhood, together with the results of our study, which has similar results with studies in the literature. We also think that further investigations are needed to evaluate its efficacy in focal and different type of seizures of childhood.

**Keywords:** *Childhood, Focal epilepsy, Lacosamide.*

## New treatment options in rare diseases

**Dr. Işıl Özer**

*Ondokuz Mayıs Un. Medical Faculty Child Metabolism Division, Samsun, Turkey*

*E-mail: isil.ozel@omu.edu.tr*

*Orcid ID: <https://orcid.org/0000-0002-3144-2915>*

### **Abstract**

The term rare disease is used for diseases with a prevalence of <2000 people in the community. For the first time in 1990, Sweden became the country that established an information center on rare diseases. Turkey has made the relevant legal arrangements in 2020. The largest group under this common roof is "Hereditary Metabolic diseases". The number of inherited metabolic diseases has reached a remarkable scope in the light of the rapidly accelerating developments in biochemistry, genetics, pharmacology, electronics sciences in 1902, when this name was first used, "Alkaptonuria" disease. In the light of the information obtained, 4 separate subgroups were created according to their common characteristics in order to produce solutions for this large group. Treatment options for metabolic disorders are both simple and complex. Diet therapy, cofactor therapy, enzyme replacement therapy (ERT), Substrate reduction therapy, chaperone therapy, tissue- organ- stem cell transplantation and gene therapy can be listed as these treatment options. Preimplantation genetics has been a rational solution to preventing disease formation, also supported by our ministry of health.

**Keywords:** *Rare diseases; Hereditary metabolic diseases; Preimplantation genetics*

# Evaluation of the Effects of CoenzymeQ10 Treatment on Corneal Epithelium by VEGF and VEGFR in STZ-induced Diabetes Rats

Dr. Çiğdem Karaca<sup>1</sup>, Assoc. Prof. Müberra Akdoğan<sup>2</sup> and  
Assoc. Prof. Ruhi Türkmen<sup>3</sup>

Afyonkarahisar Health Science University, Afyonkarahisar, Turkey <sup>1</sup>

Afyonkarahisar Health Science University, Afyonkarahisar, Turkey <sup>2</sup>

Afyonkarahisar Kocatepe University, Afyonkarahisar, Turkey <sup>3</sup>

E-mail <sup>1</sup>: drc\_karaca@hotmail.com, E-mail <sup>2</sup>: mbrakdogan@yahoo.com, E-mail <sup>3</sup>: ruhiturkmen@gmail.com

## Abstract

**Introduction:** Diabetes is a metabolic disease with microvascular damage in the chronic period. The cornea is an avascular, transparent tissue that functions as the anterior refractive surface of the eye. In the chronic period of diabetes, severe vision reduction is expected with corneal neovascularization. VEGF one of the most important mediators of angiogenesis, is upregulated during neovascularization. It has been shown in previous studies that VEGF and VEGFR expressions are increased in diabetic corneas. Coenzyme Q 10 (CoQ10) is an essential cofactor that takes electrons from complex 1 and 2 in the electron transport chain and is a molecule that has an active role in the mitochondrial electron transport chain, which has a key importance in energy production in the body. In this study, it was aimed to evaluate the effects of COQ10 administered systemically on VEGF and VEGFR immunoreactivity in corneal epithelium in rats with diabetes with STZ.

**Method:** In the study, 20 rats weighing between 200-250 g were divided into four groups as Diabetes (group 1), Diabetes + CoQ10 (group 2) only CoQ10 (group 3) and Control (group 4). Control and CoQ10 alone were administered with 0.01 M sodium citrate buffer (ph = 4.5) intraperitoneally (i.p.), other rats 50 mg / kg STZ in 0.01 M sodium citrate buffer (pH = 4.5) i.p. at once. Animals in groups 1 and 2 were diabetic by giving STZ. In groups 2 and 3, 0.35 mg CoQ10 was dissolved in 50 µl olive oil and administered by gavage for two months. At the end of 2 months, 100 mg / kg ketamine was injected to the rats and rats were sacrificed, the eyes were enucleated and placed in 10% neutral formalin. After tissue follow-up, sections were taken from the tissues embedded in paraffin, and immunohistochemistry staining protocol was applied with VEGF and VEGFR. H scoring was done by counting VEGF and VEGFR positive stained cells in microscopic evaluation.

**Results:** In our study, VEGF and VEGFR staining intensities were increased in the group 1 (p = 0.015). In the group 2, it was found to be decreased compared to the group 1 (p = 0.035). There was no difference between group 3 and group 4 (p = 0.17).

**Conclusion:** Diabetes causes damage and inflammation by increasing vascularity through the angiogenic molecules VEGF and VEGFR in the cornea. Although antiangiogenic agents and steroids are generally used for treatment, prevention of the disease is essentially the primary goal. In our study, the decrease in staining in VEGF and VEGFR in the diabetic group given CoQ10 was seen as an important marker in the reduction of corneal neovascularization and CoQ10 was considered as a reliable agent that can be used in addition to treatment.

**Keywords:** Cornea, Diabetes, CoQ10, VEGF, VEGFR

# Investigation of Colistin Resistance in Gram Negative Bacteria by Colistin Susceptibility Tube Test

**Kübra Hacıeminoglu Ülker, M.Sc<sup>1</sup>, Dr. Yeliz Tanriverdi Çaycı<sup>2</sup>  
Dr. Eda Köprü<sup>3</sup> and Dr. Asuman Birinci<sup>4</sup>**

*Ondokuz Mayıs University, Faculty of Medicine, Department of Medical Microbiology, Samsun, Turkey<sup>1</sup>*

*Ondokuz Mayıs University, Faculty of Medicine, Department of Medical Microbiology, Samsun, Turkey<sup>2</sup>*

*Ondokuz Mayıs University, Faculty of Medicine, Samsun, TURKEY<sup>3</sup>*

*Ondokuz Mayıs University, Faculty of Medicine, Department of Medical Microbiology, Samsun, Turkey<sup>4</sup>*

*E-mail<sup>1</sup>: kubra.hacieminoglu@gmail.com, E-mail<sup>2</sup>: yeliztanriverdi@gmail.com, E-mail<sup>3</sup>: edakpru@gmail.com*

*E-mail<sup>4</sup>: asumanbirinci@yahoo.com*

## Abstract

**Background:** Colistin is an important antimicrobial used as a last option in the treatment of gram-negative bacteria with multi-drug resistance and carbapenem resistance. In 2017, a joint working group of the Clinical and Laboratory Standards Institute (CLSI) and the European Committee on Antimicrobial Susceptibility Testing (EUCAST) recommended broth microdilution (BMD) as the reference method for testing colistin. However, this method is difficult and time-consuming. As a result, new methods that are simpler and faster are needed for determining colistin resistance.

**Purpose:** The purpose of this study is to compare the colistin susceptibility tube test (CSTT), which is a more effortless and simpler test to determine the colistin resistance, with the BMD.

**Methodology:** Fifty Enterobacteriales isolates that colistin resistance was determined by the reference method were included in the study. CSTT was performed with one tube containing 5 mL of Cation-Adjusted Mueller Hinton Broth and 1 colistin disk (10 µg) to give a final concentration of 2 µg/mL (test tube) and one tube without disk (growth control). After 60 minutes of incubation at room temperature for colistin disk elution, an aliquot with 25 µL of the bacterium inoculum (10<sup>8</sup> CFU/mL) was added to the tubes. Tubes were incubated for 16–20 h at 35°C and the isolate was considered resistant when bacterial growth was observed in the tube test.

**Results:** With reference method; 44 of the isolates were found colistin resistant, 6 of them were found colistin susceptible. According to the results of the CSTT method; of the 44 isolates that are colistin resistant, one isolate is colistin susceptible (2.3%), the others are colistin resistant (97.7%), of the 6 isolates that are colistin susceptible, four isolates were colistin susceptible (66.7%) and two isolates were colistin resistant (33.3%).

**Conclusion:** In our study, CSTT showed similar results to BMD in determining colistin resistant isolates. The obtained results show that a new method, CSTT is successful in determining colistin resistant isolates, and these results are promising. On the other hand, the correct determination rate of susceptible isolates is lower than the correct determination rate of resistant isolates. Therefore, the number of susceptible isolates should be increased and the study should be expanded.

**Keywords:** *Colistin Resistance, Gram Negative Bacteria, Susceptibility.*



# Early Detection of Antibiotic Susceptibility of Enterobacteriaceae Family Isolates Using an Inverted Microscope; Preliminary Study

Dr. Kemal Bilgin<sup>1</sup>, Dr. Yeliz Tanrıverdi Çaycı<sup>2</sup>, Dr. Demet Gür Vural<sup>3</sup> and Dr. Asuman Birinci<sup>4</sup>

Ondokuz Mayıs University, Faculty of Medicine, Samsun, Turkey <sup>1,2,3,4</sup>

E-mail <sup>1</sup>: kemal.bilgin@omu.edu.tr, E-mail <sup>2</sup>: yeliztanriverdi@gmail.com, E-mail <sup>3</sup>: demet.gur@yandex.com.tr,

E-mail <sup>4</sup>: asumanbirinci@yahoo.com

## Abstract

**Background:** Studies to develop new alternative methods giving results in a short time in the detection of antibiotic resistance continue to be an exciting subject.

**Purpose:** Our study aimed to determine early detection of gentamicin and ciprofloxacin antibiotic susceptibilities in Enterobacteriaceae family isolates using an Inverted microscope.

**Methodology:** In our study, 26 isolates, 17 *Escherichia coli*, and 9 *Klebsiella pneumoniae*, isolated from samples routinely sent to the Microbiology Laboratory of Ondokuz Mayıs University, Faculty of Medicine Hospital, were included. The identification of the clinical isolates included in the study was made by routine procedures applied in the Medical Microbiology Laboratory. In addition, *E. coli* ATCC 25922 isolate was also used in the study.

In our study, the determination of the minimum inhibitory concentrations (MIC) of gentamicin and ciprofloxacin antibiotics was performed by the standard microdilution method using 96-well U-bottom microplates. Ten dilutions were prepared with final concentrations of 64 - 0.125 µg/ml for gentamicin (Sigma-Aldrich) and 8 - 0.015 µg/ml for ciprofloxacin (Sigma-Aldrich). The microdilution method with the same antibiotic and concentrations was prepared with 96-well flat-bottom plates for early MIC detection by an Inverted microscope (Olympus). Plates were stored at -20°C until use.

The prepared inoculum was placed on both plates. After the inoculation, the U-bottom microplates were incubated at 36°C overnight. Flat bottom microplates were incubated for five hours at 36°C. Then, we tried to determine the MIC according to the growth present in the wells by examining with an Inverted microscope.

**Results:** The comparison of standard microdilution and flat plate microdilution results of both antibiotics for all isolates included in the study is given in the Table.

**Table 1.** Compatibility rates of MIC results obtained from flat plate microdilution and standard microdilution methods.

		≥-5 (%)	-4 (%)	-3 (%)	-2 (%)	-1 (%)	0 (%)	+1 (%)	+2 (%)	+3 (%)	+4 (%)	≥+5 (%)
<i>E. coli</i> (18)	Gentamicin		1 (5.6)			4 (22.2)	9 (50)	4 (22.2)				
	Ciprofloxacin					2 (11.1)	15 (83.3)					1 (5.6)
<i>K. pneumoniae</i> (9)	Gentamicin						8	1				
	Ciprofloxacin						8		1			

**Conclusion:** In *E. coli* isolates, for gentamicin, 9 isolates fully compatible, 4 isolates with -1 dilution difference and 4 isolates with +1 dilution difference; For ciprofloxacin, 15 isolates fully compatible and 2 isolates with -1 dilution difference were detected. In *K. pneumoniae* isolates, for gentamicin, 8 isolates fully compatible, and 1 isolates with +1 dilution difference; For ciprofloxacin, 8 isolates fully compatible were determined. The method's applicability is not yet clear due to the low number of microorganisms included in our preliminary study. Therefore, to understand the applicability of the method, comparative new studies with standard methods using a large number of Enterobacteriaceae isolates with the use of different antibiotics are needed.

**Keywords:** *E. coli*; *K. pneumoniae*; MIC; Inverted microscope

# Optimization Study of Detecting Antibiotic Sensitivity of *Pseudomonas aeruginosa* Using an Inverted Microscope

**Dr. Kemal Bilgin**

Ondokuz Mayıs University, Faculty of Medicine, Samsun, Turkey  
E-mail: kemal.bilgin@omu.edu.tr

## Abstract

**Background:** Although there are standard methods for detecting antibiotic resistance, studies continue on alternative methods that resulted in a shorter time.

**Purpose:** Our study investigated the detection hour of the antibiotic sensitivity of *Pseudomonas aeruginosa* to gentamicin and ciprofloxacin by using an inverted microscope.

**Methodology:** *P. aeruginosa* ATCC 27853 standard strain was used as an isolate in our study. In our study, the determination of the minimum inhibitory concentrations (MIC) of gentamicin (Sigma-Aldrich) and ciprofloxacin (Sigma-Aldrich) antibiotics was done by microdilution method using 96-well U-bottom microplates. Ten dilutions were prepared with final concentrations of 64 - 0.125 µg/ml for gentamicin and 8 - 0.015 µg/ml for ciprofloxacin. The microdilution method was prepared in 96-well flat-bottom plates, with the same antibiotic and concentrations for optimization trials of early MIC detection by Inverted microscope (Olympus).

The prepared inoculum was applied in the same shape and amount on both plates. After the inoculation, the U-bottom microplates were incubated at 36°C overnight. Flat bottom microplates were incubated at 36°C, and the plates were examined with an Inverted microscope every hour for eight hours, and MIC determination was attempted according to the presence of growth in the wells. Then the flat bottom microplates were incubated overnight, and the MIC value was determined depending on the color change due to the pigments created by *P. aeruginosa*.

**Results:** The flat plate results were compared with the standard microdilution results and determined as equal, one-fold up or down concentrations.

**Table1.** Comparison of MIC results by microdilution and inverted microscope.

Trial	Antibiotic	MIC	Hour								
			1	2	3	4	5	6	7	8	24*
1	Gentamycin	2	-	-	-	2	1	1	1	1	2
	Ciprofloxacin	0.25	-	-	-	0,5	0,5	0,25	0,5	0,25	0,5
2	Gentamycin	2	-	-	-	2	1	1	1	1	2
	Ciprofloxacin	0,25	-	-	-	0,25	0,25	0,25	0,25	0,25	0,25
3	Gentamycin	2	-	-	-	-	1	1	1	1	2
	Ciprofloxacin	0,25	-	-	-	0,25	0,25	0,25	0,25	0,25	0,25

\*: MIC determination by color change due to pigment formation.

**Conclusion:** In our study, no bacteria were found in the wells for the first three hours. Bacteria could be seen at the bottom of the wells at the fourth hour, except for the gentamicin antibiotic of the third trial. However, a clear image could be taken at the seventh and eighth hours. The method's applicability is not clear yet, since clinical isolates were not included in our optimization study. In order to understand the applicability of the method, comparative studies with standard methods using

a large number of *P. aeruginosa* isolates obtained from clinical samples with the use of different antibiotics are needed.

**Keywords:** *Pseudomonas; aeruginosa; MIC; Inverted microscope*

# The distribution of Enterobacteriales isolated from the urine samples of children, and the evaluation of antimicrobial susceptibility

Dr. Yeliz Tanrıverdi Çaycı<sup>1</sup>, Dr. Gülşah Karacan<sup>2</sup>, Dr. Moein Yoosefi<sup>3</sup>,  
Dr. Kemal Bilgin<sup>4</sup>, Dr. Demet Gür Vural<sup>5</sup> and Dr. Asuman Birinci<sup>6</sup>

Ondokuz Mayıs University, Faculty of Medicine, Samsun, Turkey

E-mail<sup>1</sup>: yeliztanriverdi@gmail.com, E-mail<sup>2</sup>: gulskkaracan29@gmail.com, E-mail<sup>3</sup>: moeinyoosefi17@gmail.com,  
E-mail<sup>4</sup>: kemal.bilgin@omu.edu.tr, E-mail<sup>5</sup>: demet.gur@yandex.com, E-mail<sup>6</sup>: asumanbirinci@yahoo.com

## Abstract

**Background:** Antimicrobial resistance has become a serious problem as a result of extensive antibiotic consumption.

**Purpose:** In this study we aimed to determine the bacteria grown in urine cultures of children and antimicrobial resistance rates in our region.

**Methodology:** The results of urine cultures that sent from Department of Pediatrics between 01.01.2015 and 31.12.2020 to the Microbiology Laboratory, were evaluated retrospectively. The identification of the bacterial growth was tested in Vitek MS (Biomeriux, France) and antimicrobial susceptibility of the isolates were tested in Vitek2 Kompakt (Biomeriux, France).

**Results:** Total of 5511 urine culture samples were evaluated. The most frequently isolated bacteria was *E. coli* 67.5% and followed by *Klebsiella* spp 21.1%, *Proteus* spp 4.81%. The highest resistance rates were determined against ampicillin in *E. coli* and *Klebsiella* spp isolates (78.96%, 99.91%). The lowest resistance rate was determined against imipenem and meropenem for *E. coli* (0.12%, 0.15%) isolates and amikacin and nitrofronatin in *Klebsiella* spp isolates (6.10%, 2.46%).

**Conclusion:** Order to provide effective treatment in urinary tract infections and to minimize morbidity and long-term complications, the frequency of the pathogens causing urinary tract infections, local or regional antibiotic resistance should be known and empirical treatment should be arranged accordingly.

**Table 1.** The antimicrobial resistance rate of *Enterobacteriaceae* isolates.

	<i>E. coli</i>	<i>Klebsiella</i>	<i>Enterobacter</i>	<i>Proteus</i>	<i>Citrobacter</i>	<i>Serratia</i>	<i>Providencia</i>	<i>Morganella</i>	<i>Raotuel la</i>	<i>Salmonella</i>
<b>Ampicilin</b>	78.96	99.91	99.96	40.28	100	100	100	100	100	20
<b>Amoxisilin/ clavulonika cid</b>	15.21	19.34	39.21	3.80	20.48	25.04	0	33.75	0	0
<b>Cefuroxim</b>	42.96	63.11	54.06	13.30	33.28	81.38	66.68	82.50	0	0
<b>Cefepim</b>	5.94	14.18	8.16	4.56	10.24	0	0	7.50	0	0
<b>Ceftriaxone</b>	39.45	60.96	55.08	9.12	30.72	15.65	50	41.25	0	0
<b>Ceftazidim</b>	35.64	53.74	52.53	8.36	28.16	12.52	66.68	37.50	0	0
<b>Ciprofloxac in</b>	23.85	27.94	10.71	11.40	12.80	0	50	27.50	0	20
<b>Ertapenem</b>	1.17	19.51	17.34	3.42	7.68	9.39	50	1.25	0	0

<b>Imipenem</b>	0.12	6.44	6.12	0.38	2.56	0	33.34	1.25	0	0
<b>Meropenem</b>	0.15	7.30	5.61	0	0	0	33.34	0	0	0
<b>Amikacin</b>	1.11	6.10	3.57	1.90	5.12	0	16.67	0	0	0
<b>Gentamicin</b>	13.98	33.10	23.46	14.44	12.86	12.52	33.34	28.75	0	0
<b>Fosfomisin</b>	1.50	21.32	35.19	15.58	0	6.26	33.34	97.50	66.66	0
<b>Nitrofurantoin</b>	4.17	2.46	4.59	79.80	2.56	75.12	100	75	0	20
<b>Trimetoprim/sulfametaxazol</b>	51.36	55.97	31.11	45.60	25.60	9.39	50	33.75	0	20

**Keywords:** *The globalization of science; scientific collaboration; globalizing knowledge economy*

## Future Expectations in ART

Assoc. Prof. İlay Gözükara<sup>1</sup> and Prof. M. Turan Çetin<sup>2</sup>

Private Prof.Dr.Turan Cetin IVF Centre, Turkey<sup>1,2</sup>  
E-mail <sup>1</sup>: ilayozt@gmail.com, E-mail <sup>2</sup>: mtcetin@yahoo.com

### Abstract

There are some troubles during In Vitro Fertilisation (IVF) treatment even though advances. Embryo selection, advanced maternal age and fertility preservation are most common ones. Firstly, the selection of viable embryo remains a principle challenge of IVF in order to predict live birth. Artificial intelligence could provide benefit in the future. Dataset based on time-lapse images combined with embryo characteristics may provide the selection of accurate embryo. Left over products of IVF for their contents in proteins, metabolites, nucleic acid and -omics technologies may also help embryo selection and growing that will likely change the current IVF procedures. The laboratory environment may be completely identical to the in vivo environment, and as a result, the fetus will be grown in vitro until birth in the future. Secondly advanced maternal age is a critical clinical issue in IVF cycles. Emerging therapeutic approaches have been proposed for these patients. Germ line engineering could be considered to restore the competence of aged oocytes. Chromosome therapy is another promising one. In the future such practice might find direct application in the treatment of chromosomal disorders but might also apply to the correction of aneuploid germ cells and embryos in IVF. Beside chromosome therapy it may be possible to generate new gamet cells from stem cell in the future. It was already reported that murine induced pluripotent stem cells were differentiated into functional oocytes in the presence of specific growth-factors. Thirdly some future fertility preservation methods can be needed like some new gonadoprotective agents, also some new drugs to improve tissue grafting. Artificial ovaries containing isolated follicles and culturing primordial follicles to mature oocyte might serve as a revolutionary method without the danger of cancer reseeding. In conclusion we may not see any pregnant women in the future even may not see any children with syndrome and congenital diseases in the future because of some advanced technologies.

**Keywords:** *Germ line engineering; chromosome therapy; IVF; gamet cells.*

# Can We Consider Embryos Solely as Biological Material?

**Maide Barış**

*Istanbul Health and Technology University, Department of Medical History & Ethics, Faculty of Medicine, Istanbul, Turkey  
E-mail: baris.maide@gmail.com*

## **Abstract**

Traditionally, bioethical discussions about embryo research tend to resort to moral status arguments. Pro-embryo research argumentators in bioethics who embrace what we call “none-approach” use “psychological arguments” or “physiological arguments” to claim that embryos do not have any moral status. On the other hand, major anti-embryo research argumentators usually adopt what we call “all-approach” use “development-based moral arguments” and “metaphysical moral arguments” to defend that embryos have moral status. This paper firstly presents these moral approaches briefly, maintaining that the “moral status” the approach does not help in practice, especially regarding interventions in preimplantation-stage embryos in the laboratory. While embracing and respecting great moral diversity of beliefs about the embryo, this paper suggests that there should be a practice-focused and less counter-intuitive moral stance somewhere in between “embryos are complete humans who have the full moral status” and “embryos are nothing but accumulations of cells.” Secondly, we will argue that although embryos are not entitled to full moral status as much as a paradigm human does, they are still worthy of moral respect and cannot be considered solely as biological materials because (i) they have inherent potential to become one of us; (ii) they are human organisms in a biological sense and (iii) they are soon to be considered as patients in near future, rather than only as biological research entities, at a time when Nobel-winning CRISPR/Cas9-based germline genetic intervention is waiting at the doors of reproductive clinics. Finally, we will maintain that the statement “embryos are worthy of moral respect” does not say anything about the post-implantation stage, especially regarding abortion. It only contends that because embryos are worthy of moral respect and to be considered patients and subjected to genetic intervention in a petri dish, then they need to be handled by a physician/ specialist who is trained microsurgery to deal with embryos: A clinical embryologist.

**Keywords:** *Embryo Research; Moral Status; Moral Respect; Bioethics.*



# Common Chest CT Findings in 100 COVID-19 Patients Followed for Pneumonia

**Dr. Mehmet Akçiçek**

*Malatya Turgut Özal University, Faculty of Medicine, Malatya Training and Research Hospital, Radiology Department,  
Malatya/Turkey  
E-mail: mmakcicek@gmail.com*

## Abstract

**Background:** CT has become an important imaging method in the diagnosis and treatment of patients with viral pneumonia. The use of low-dose CT scanning for the diagnosis of viral pneumonia optimizes patient management by providing rapid treatment of patients suspected of COVID-19 infection. As shown in previous studies; chest CT images may show different imaging features in COVID-19 patients depending on the period and severity of the disease.

**Purpose:** In this study, we investigated the common CT imaging features and changes of COVID-19 in 100 pneumonia patients, who were confirmed by RT-PCR and underwent control CT examination during follow-up.

**Method:** The patients who admitted in July, August and September 2020 when the highest rate of patient admission and hospitalization in our hospital during the COVID-19 pandemic and whose RT-PCR test positivity was confirmed has been scanned retrospectively. Patients whom was detected pneumonia in chest CT examined at their first visit and a second CT performed in the 5-14 day period in their follow-up, were included. A total of 100 patients, 61 of whom are male, between the ages of 25-90 were identified. CT examinations were performed on 16 and 128 MDCT devices in low dose. Images were transferred to the workstation and analyzed in the PACS system. The findings of the first and second CT of each patient were defined according to the Fleischner community and literature information.

**Results:** The most common findings in the first CT were GGO (Ground glass opacity) (68%), consolidation (46%), nodule (31%) and reticulation (28%). Findings other than these are as follows; airway changes (17%), vascular enlargement (12%), subpleural line (8%), halo (2%), reverse halo (2%), pleural effusion. 94% of the lesions were peripheral and 92% were multifocal. In the control CT examinations performed within 4-15 days, the most common findings were GGO (70%), consolidation (70%), 65% reticulation, 31% CPP, 33% airway changes. Vascular enlargement in 27%, subpleural line in 22%, subpleural fibrotic strip in 13%, nodule in 13%, pleural effusion in 6%, reverse halo findings in 4%, and pericardial effusion in 1% of the patients were found. Pleural thickening (1%) and lymphadenopathy (11%), which were also detected in the first examinations were detected. 78% of the lesions were peripheral, 22% were peripheral and centrally located, and 99% were multifocal.

**Conclusion:** CT findings of COVID-19 pneumonia resemble those of many other infectious and non-infectious diseases, especially viral pneumonia types. These findings may vary depending on the disease phase during follow-up, individual characteristics of the patient, underlying diseases and treatment interventions.

**Keywords:** COVID-19, Chest, CT, Pneumonia

# Does COVID-19 infection trigger the formation of Anti-nuclear Antibodies?

Dr. Melek Bilgin<sup>1</sup> and Dr. Eşe başbulut<sup>2</sup>

Health Sciences University, Samsun, Training and Research Hospital, Medical Microbiology Department, Samsun, Turkey <sup>1,2</sup>

E-mail <sup>1</sup>: drmelekbilgin@gmail.com, E-mail <sup>2</sup>: Ese.Basbulut@saglik.gov.tr

## Abstract

**Aim:** Increasing evidence suggests that autoimmunity may play a role in the pathophysiology of SARS-CoV-2 infection during both the acute and ‘long COVID’ phases of disease. However the long-term consequences of COVID-19 and whether it will trigger autoimmunity in the future are curious and require further research. There is insufficient knowledge on the assessment of autoimmune antibodies in convalescent SARS-CoV-2 patients. The aim of this study is to evaluate the presence of antinuclear antibodies (ANAs) in convalescent SARS-CoV-2 patients and to compare with ANA results before the onset of COVID-19.

**Material and Methods:** 23 healthcare workers with Covid 19 infection (SARS CoV-2 PCR test positive) whose ANA test results were known and reported before the Covid 19 infection were included in this study. ANA tests were studied from the serum of participants at least 3 months after Covid-19 infection.

Diagnosis of COVID-19 infection was performed by a RT-qPCR method. Bio-Speedy, SARSCoV-2 Double Gene RT-qPCR Detection Kit (Bioeksen, İstanbul, Türkiye) were used for detection of virus RNA. ANA positivity and patterns were assessed by indirect immune-fluorescence on Hep-2 cells (Euroimmun, Luebeck, Germany) using standard techniques.

**Results:** The study included 23 people, 19 women and four men, with a mean age of 45. Differences were found in the ANA IFA pattern in 3 of 23 patients after Covid-19 infection compared to pre-infection. In the first case, the ANA IFA pattern before Covid-19 was speckled weakly positive, while the ANA IFA result was detected as anti-nuclear membrane positive (++) 3 months after infection. Thereupon, anti-mitochondrial antibody (AMA), anti-smooth muscle antibody (ASMA), and anti-LKM were studied for autoimmune liver disease in the patient and found as negative. Also, RNP / Sm, Sm, SS-A, Ro 52, SS-B, Scl-70, PM-Scl, Jo-1, CENP B, PCNA, dsDNA, nucleosomes, histones, rib P-prot., AMA-M2, and DFS70 antibodies were studied by immunoblot method, and anti-RNP / Sm were detected positive. While the second case was chromosomal granular (GGK) positive before infection, ANA IFA was found homogeneous positive, and cytoplasm speckled positive after infection. Then Immunoblot ANA wide panel was studied in the case, and Scl-70, DFS70, dsDNA found positive. In the third case, while the previous result was negative, it was determined as granular weak positive after infection. In this case Ena blot result was evaluated as negative. One year before they had the Covid-19 infection, all 3 cases had negative Ena blot results.

**Conclusion:** As a conclusion, our study revealed that Covid-19 infection can trigger autoimmunity, and patients should be followed up for a long time after Covid-19 infection. Further studies with larger patient series are needed to better define the clinical significance of auto-antibody positivity and to understand the possible role of Covid-19 in autoimmune diseases.

**Keywords:** SARS-CoV-2; COVID-19; Anti-nuclear Antibodies; Autoimmunity

# Guillain Barre Syndrome Induced by Covid 19: Discussion with a Case

**Süsen Banazılı, MD<sup>1</sup> and Asst. Prof. Mesut Öterkus<sup>2</sup>**

*Malatya Training and Research Hospital, Anesthesiology and Reanimation Clinic <sup>1</sup>  
Malatya Turgut Özal University Faculty of Medicine Department of Anesthesiology and Reanimation <sup>2</sup>  
E-mail <sup>1</sup>: drsusen@hotmail.com, E-mail <sup>2</sup>: mesutoterkus@hotmail.com*

## Abstract

Covid 19, which is among the most important health problems of today, affects many organs and systems. The nervous system is one of them. Many complications such as stroke, encephalitis, necrotizing encephalitis, and Guillain Barre syndrome (GBS) have been reported in the literature. GBS is an autoimmune-induced demyelinating polyneuropathy. It usually occurs following a recent infection. Covid 19 is one of these infections. The main mechanism is thought to be T cell activation or similarity of virus proteins to nerve cell protein. The most effective treatments are immunotherapy (plasma exchange, IVig) and physiotherapy. It leaves permanent neurological sequelae in approximately 20% of patients.

We aimed to present a 51-year-old male patient who developed GBS. Our patient did not have a recent history of any infection or neurological deficit other than covid 19 infection. Our patient was intubated electively due to increased respiratory distress and with the improvement of respiratory parameters, the patient was extubated decreased oxygen saturation during his follow-up and treatment in the intensive care unit. The patient was extubated as the respiratory parameters improved after the treatment. In his examination after extubation, bilateral lower extremity and loss of strength and reflexes in the left upper extremity were observed. GBS was considered in the foreground. Since the EMG was also compatible, IVIG (0.4g / kg) treatment was started for 5 days. Physiotherapy was started on the patient in consultation with a physical therapy and rehabilitation specialist. After 4 weeks, the patient started walking with support. In literature,

GBS after covid 19 is usually a case report. In a study conducted in Italy; showed that GBS increased 2.6 times during the pandemic period of covid 19. As a result, It should be kept in mind that GBS can also appear as a Covid 19 complication and early treatment increases the chance of success.

**Keywords:** Covid-19, Autoimmunity, Guillain Barre syndrome

# Clinical experience in patients with Adult Still's disease

**Dr. Muhammed Okuyucu**

*Ondokuz Mayıs University Faculty of Medicine, Department of Internal Medicine, Samsun, Turkey  
E-mail: muhammedokuyucu55@gmail.com*

## Abstract

**Background:** Adult Still's disease (ASD) is an inflammatory disease of unknown etiology which is characterized by daily (quotidian) fever attacks, arthritis and evanescent rash. The most typical symptom of the disease is the non-itchy pink rash that occurs especially on the body, arms, and legs accompanied by fever attacks. Other symptoms of the disease are sore throat, swelling of lymph nodes, enlargement of the liver and spleen, and elevated liver test results. Although there is no specific laboratory test used in the diagnosis of the disease, the most important test used is ferritin, which is high in 70% of the patients. Steroid and methotrexate treatments are mostly sufficient in the patients. Other immunosuppressive and anti-TNF agents are used in patients who do not respond adequately or cannot tolerate treatment.

**Purpose:** In this study, we examined our 7 years of experience in ADS in the rheumatology clinic of Ondokuz Mayıs University (OMU).

**Methodology:** 54 patients who received the diagnosis of ASD and were followed up in OMU rheumatology clinic between February 2014 and March 2021 were retrospectively analysed in terms of clinical findings, involvement, laboratory results, and treatment.

**Results:** Data of 54 patients were compiled in total. 61.1% of the participants are women (n=33); the general age average is  $40.17 \pm 16.8$ . The most common symptom is fever (96.3%, n = 52), the most commonly administered medication is steroid (100%, n = 54) (Tables 1 and 2). According to their current status, 9.3% (n = 5) of the patients died, 20.4% (n = 11) of them were in remission, and 70.4% (n = 38) of them are still being treated. Those who develop Macrophage Activation Syndrome (MAS) are less likely to be in remission (p=0.013). Types of medication and laboratory data do not affect the current status of the patients (p> 0.05).

**Table 1.** Clinical findings

Finding	%	n
Fever	96.3	52
Arthritis	92.6	50
Rash	63.0	34
Sore throat	50	27
LAP	35.2	19
Hepatomegaly	16.7	9
Splenomegaly	14.8	8
MAS	9.3	5
Cardiac involvement	5.6	3
Lung involvement	3.7	2
Hemophagocytic syndrome	3.7	2

**Table 2.** Medications administered

Medication	%	n
Steroid	100	54
Methotrexate	94.4	51
Tocilizumab	35.2	19
Cyclosporine	11.1	6
Leflunomide	11.1	6
Anakinra	3.7	2
Cyclophosphamine	1.9	1
Infliximab	1.9	1

**Conclusion:** ASD is one of the major diseases that should come to the mind of clinicians in the differential diagnosis of patients with fever, arthritis, and rash in particular. Clinicians should pay attention to MAS because it decreases the rate of remission and causes serious clinical disorders in patients followed up due to this diagnosis.

**Keywords:** *Adult Still's disease, clinical experience, patients*

## Distal metatarsal Chevron osteotomy + modified McBride method in hallux valgus, short term 1-3 year results

**Dr. Muhammet Salih Ayas**

*Republic of Turkey Ministry Of Health, Erzurum Regional Education and Research Hospital, Erzurum/Turkey  
E-mail: muhammetsalihayas@yahoo.com.tr*

### Abstract

**Background:** Short-term efficacy of distal metatarsal Chevron osteotomy + modified McBride method in hallux valgus.

**Purpose:** In this study, the results of distal metatarsal Chevron osteotomy + modified McBride method in patients with a diagnosis of mild disease (HVA  $\leq$  40, IMA  $<$ 13) hallux valgus were evaluated.

**Methodology:** Twenty-six feet of 22 patients with hallux valgus deformity, who were operated between 2017 and 2020 in Erzurum Regional Training and Research Hospital, were operated with distal metatarsal Chevron osteotomy + modified McBride method. Radiological, functional and pain evaluations were examined.

**Results:** Preoperative and postoperative hallux valgus angle and 1.-2. There was a statistically significant difference between the intermetatarsal angle values ( $p < 0.001$  for both values), and no statistically significant difference was found between the 1st distal metatarsal articular angle values ( $p > 0.5$ ). The average visual analog scale values, which were  $5.6 \pm 1.0$  before the operation, decreased to  $1.4 \pm 1.0$  after the follow-up and this result was found statistically ( $p < 0.001$ ). When the results of the control American Orthopedic Foot-Ankle Association Score ( $86.2 \pm 8.2$ ) were compared with the preoperative values ( $34.2 \pm 14.8$ ), a statistically significant difference was found ( $p < 0.001$ ).

**Conclusion:** In conclusion, distal metatarsal Chevron osteotomy + modified McBride operation is an effective and safe technique in the surgical treatment of hallux valgus. Successful results have been demonstrated in reducing pain and correcting radiological and cosmetic deformity.

**Keywords:** *Hallux valgus, Chevron, Modified McBride, Osteotom*

# Effects of Avanafil, A Phosphodiesterase Type 5 (Pde-5) Inhibitor, on A549 Lung Cancer Cell Proliferation

**Asst. Prof. Muhammet Çelik**

*Atatürk University, Faculty of Medicine, Department of Medical Biochemistry, Erzurum, Turkey  
E-mail: drmuhammetcelik@gmail.com*

## Abstract

**Objective:** Cyclic nucleotide phosphodiesterase enzymes (PDEs) are a family of super enzymes found in all cell types and tissues. Different enzymes are found in this enzyme family, from one to eleven (PDE1-PDE11). The functions of PDE enzymes found in different tissues of the body are, of course, different. The contribution of PDE iso-enzymes to metabolism and their roles in physiopathological events has attracted attention. These enzymes play a role in the pathophysiology of many diseases such as inflammation, neurodegeneration, cancer, immunomodulation. With the studies in recent years, new PDE inhibitors have been introduced to the market. Studies have shown the significant effects of sildenafil, a PDE-5 inhibitor, especially on lung cancer. Many different mechanisms play a role in the anticancer activity of PDE-5 inhibitors. Avanafil is also a PDE-5 inhibitor, and its effect is much higher than sildenafil. In our study, the effect of Avanafil on the viability of A549 lung cancer cells was investigated.

**Methods:** We investigated the dose and time-dependent effects of Avanafil on cell proliferation in A549 cells. For this aim, different concentrations of Avanafil (from 1  $\mu\text{M}$  to 128  $\mu\text{M}$ ) were added onto A549 cells, and viability and proliferation were evaluated with MTT test at 24 and 48 hours after incubation.

**Results:** Avanafil has been studied with concentrations of 1-2-4-8-16-32-64 and 128  $\mu\text{M}$ . When the cell viability tests were examined at 24th and 48th hours, it was observed that all doses of Avanafil reduced cell division in 24 hours, and the best effect was at 128  $\mu\text{M}$  concentration of Avanafil. When the 48th hour was examined, it was shown that the effect continued, especially the anticancer activity of Avanafil at 128  $\mu\text{M}$  concentration remained very high and significantly reduced cancer cell proliferation.

**Conclusion:** Our findings showed that Avanafil, like other PDE5 inhibitors, inhibited the proliferation in cancer cells. Studies have shown that PDE5 is expressed in colon, prostate, breast, and pancreatic cancers. These results show that PDE5 enzymes are effective in carcinogenesis. PDE5 enzyme inhibition has become an important target in cancer prevention and treatment. Intracellular apoptosis and necrosis pathways induced by PDE5 inhibition may be responsible for the anticancer action mechanism of Avanafil. However, additional laboratory studies are needed to elucidate these pathways. Our study concluded that Avanafil emerges as a new therapeutic target in lung cancer patients, although it is outside of the current indications for use. The shorter duration of action compared to sildenafil will also reduce its unwanted side effects and facilitate dose control in cancer treatment.

**Keywords:** Lung cancer; Phosphodiesterase enzymes; PDE-5; Avanafil; A549 cell line.

# Comparison of posterolateral approach and direct reduction posterior-anterior screw fixation method versus indirect reduction anterior-posterior screw fixation method for posterior malleolus fixation in ankle trimalleolar fractures

Muhammet Kalkışım, MD<sup>1</sup> and Kerim Öner, PhD, MD<sup>2</sup>

Surgeon, Department of Orthopaedic Surgery, Sorgun Public Hospital, Yozgat, Turkey <sup>1</sup>

Assistant Professor, Department of Orthopaedic Surgery, Karadeniz Technical University School of Medicine, Trabzon, Turkey <sup>2</sup>

E-mail <sup>1</sup>: muhammetkalkisim@gmail.com, E-mail <sup>2</sup>: dr.kerimoner@hotmail.com

## Abstract

**Background:** A consensus on how posterior malleolar fragments in trimalleolar fractures should be reduced and fixated still lacks. Anatomic reduction and rigid fixation are important in the treatment of posterior fragment in trimalleolar fractures.

**Purpose:** In this study we aimed to compare direct reduction (DR) using posterolateral approach and the posterior-anterior (P-A) screw fixation method versus indirect reduction (IR) with anterior-posterior (A-P) screw fixation method.

**Methodology:** In our study, patients with ankle trimalleolar fracture were evaluated retrospectively. This comparative study reviewed a case series from 2017 to 2020. Patients with at least 1 year postoperative follow-up were included in the study. Patients were categorized into two groups. The first group consisted of patients who underwent fixation with direct reduction (DR) and PA screws, and the second group consisted of patients who underwent fixation with indirect reduction (IR) and AP screws. Demographic datas, American Ankle Society scores (AOFAS), ankle range of motions (ROM), fracture types (according to Lauge-Hansen classification), the size of posterior malleolar fragment, complications and functional bone healing time were compared in both groups. Postoperative X-rays were performed to determine the residual step-off in the articular reduction. The degree of osteoarthritis was evaluated by final follow up controls with x-rays.

**Results:** 16 patients (9 males, 7 females) in group DR and 12 patients (6 males, 6 females) in IR group were recruited. The mean age in group DR were 41.6 (20-56) while 37.1 (21-53) were in group IR. There was no significant difference between the two groups in terms of complication rates, functional bone healing times, follow-up times, fracture types, amount of intra-articular step-off and ROM ( $p > 0.05$ ). The AOFAS scores were significantly better in the IR group compared to the DR group ( $p > 0.05$ ). None of the complications such as fixation failure, infection, delayed union were observed in both groups.

**Conclusion:** The results are satisfied both in two different approaches for fixation the posterior malleolus in trimalleolar fractures. However, functional results are better in the indirect reduction (IR group) method. We attribute the better functional results in indirect reduction and anterior-posterior screw fixation method because of the less invasive approach.

**Keywords:** *Posterior malleolar fractures, Trimalleolar fractures, Posterolateral approach, Screw fixation*

## Covid-19 and Comorbidities

**Dr. Mukadder Erdem<sup>1</sup>, Recai Aci<sup>2</sup>, Assoc. Prof. Dr. Mahcube Çubukçu<sup>3</sup>, Adem Keskin, PhD<sup>4</sup>, Dr. Eda Türe<sup>5</sup> and Dr. Ebru Ulaş<sup>6</sup>**

*Samsun Training and Research Hospital Biochemistry<sup>1,2,6</sup>  
Samsun Training and Research Hospital Family Medicine<sup>3,5</sup>  
Aydın Adnan Menderes University Biochemistry<sup>4</sup>*

*E-mail<sup>1</sup>: mukadder.erdem@saglik.gov.tr, E-mail<sup>2</sup>: recai.aci@saglik.gov.tr, E-mail<sup>3</sup>: mahcube@gmail.com,  
E-mail<sup>4</sup>: adem.keskin@saglik.gov.tr, E-mail<sup>5</sup>: eda.ture@hotmail.com, E-mail<sup>6</sup>: drebruu@yahoo.com*

### Abstract

**Objectives:** The pandemic situation with the emergence of severe acute respiratory syndrome coronavirus 2 (SARS-CoV-2) from China has endangered human lives. Corona virus disease 2019 (COVID-19) is presented with asymptomatic, mild, or severe pneumonia-like symptoms. COVID-19 patients with diabetes, chronic obstructive pulmonary disease (COPD), cardiovascular diseases (CVD), hypertension, malignancies, HIV, and other comorbidities could develop a life-threatening situation. In this study, the files of patients with COVID-19(+) who were followed up in the service and intensive care unit were retrospectively scanned, and their comorbid disease status was examined. This study aimed to compare with some biochemical and hematological parameters in terms of the mortality rates, taking into account the length of stay in the hospital of Covid 19 patients with comorbid and non-comorbid.

**Methods:** A total of 44 patients with COVID-19 in the intensive care unit and 71 patients in service were included in the study. We performed correlation analysis between whole blood parameters, important biomarkers for prognosis of the disease in the inflammatory process and hospitalization day, and mortality. We also evaluated the effects of these parameters of comorbidity diseases.

**Results:** When looking at the correlation of biochemical and hematological parameters of all COVID-19 (+) patients hospitalized in both the service and the intensive care unit with the day of hospitalization; it was positively correlated with PT, INR, CRP, FERRITINE, PCT, ESR, D-DIMER, WBC, NEUTROPHIL, MLR, and NLR parameters of the day of hospitalization, while it was negatively correlated with HGB and LYMPHOCYTE parameters ( $p < 0.01$ ). In addition, Ex/discharge status was positively correlated with LNF ( $p < 0.01$ ), PLT ( $p < 0.01$ ) parameters, while it was negatively correlated with PT ( $p < 0.05$ ), INR ( $p < 0.05$ ), CRP ( $p < 0.01$ ), Ferritin ( $p < 0.01$ ), PCT ( $p < 0.01$ ), ESR ( $p < 0.01$ ), D-Dimer ( $p < 0.01$ ), WBC ( $p < 0.01$ ), NTR ( $p < 0.01$ ), MLR ( $p < 0.01$ ), NLR ( $p < 0.01$ ) and Troponin-I ( $p < 0.01$ ) parameters as shown. Also, out of 115 COVID-19 (+) patients participating in this study, 91 comorbid diseases were detected, while 24 of them did not. This comorbid disease was detected in 22 patients with DM (24.18%), 21 with hypertension (23.08%), 11 with CVD (12.09%), 9 with CRF(Chronic Renal Failure) (9.89%), 6 in COPD (6.59%), 15 of them had pneumonia (16.48%), and 7 had cancer (7.69%). Besides, out of 115 COVID-19 (+) patients participating in this study, all 28 patients who died were found to have comorbid diseases. It was determined that DMD in 5 (17.86%), hypertension in 4 (14.29%), CVD in 6 (21.43%), CRF in 3 (10.71%), COPD in 2 (%7.14), Pneumonia in 4 (14.29%), cancer in 4 (14.29%) patients with Exitus.

**Conclusion:** It has been determined that whole blood and essential parameters in the inflammatory process correlated with the length of stay and mortality, these parameter levels varied in COVID-19 patients with comorbidity and lead to an increase in the size of stay and mortality rates.

**Keywords:** Covid-19, Comorbidity, Hospitalization day, Mortality.



# A Rare Form of Liver Failure and Immune Deficiency: IFNGR1 gene Mutation

**Murat Cag, PhD, MD**

*Strasbourg University, Hepatopancreatic Biliary Surgery Unit, France  
Memorial Bahcelievler Hospital, General Surgery Department, Istanbul Turkey  
E-mail: cagmurat@hotmail.com*

## **Abstract**

Immunodeficiency-27A is a disorder caused by mutations in the IFNGR1 gene that is inherited autosomal recessively. Patients with complete IFNGR1 deficiency have a severe clinical phenotype characterized by early and often fatal mycobacterial infections. Thus, the disorder can be categorized as a form of mendelian susceptibility to mycobacterial disease (MSMD). Resistant mycobacterial infections can begin from childhood. Salmonella infections and diarrhea can be seen in 5% of the patients. Index case applied to our clinic as the 3rd living child of 3rd degree relatives parents. It was learned that there was no other affected individual in the family. From her history, it was learned that there was no significant disease in the prenatal period and until his second year of age, but then gradually swelling in the abdomen, uncontrollable diarrhea, weight loss, and she had received 4-way tuberculosis treatment for the last 3 months. Laboratory tests revealed impaired liver function, high bilirubin levels and pancytopenia, very low total protein, albumin, Na and Mg values, and high CRP. In the abdominal ultrasound, liver and spleen were larger than normal and heterogeneous, and congestion in the portal circulation was observed. This mutation is known as Immunodeficiency Type 27 A and is one of the ultra-rare forms of immunodeficiency. As in our patient, resistant mycobacterial and Salmonella infections are usually seen and are usually fatal. This Syndrome causing liver and immunodeficiency was presented as an ultra rare Syndromic case.

**Keywords:** *Liver failure, immune deficiency, IFNGR1, whole exome sequencing, WES*

# Commitment to the medical profession and individual competitiveness of Medical Faculty students in Turkey

**Asst. Prof. Mustafa Bayraktar**

*Atatürk University, Faculty of Medicine, Department of Family Medicine, Erzurum, Turkey  
E-mail: mustafabayraktar@atauni.edu.tr*

## Abstract

**Objective:** To investigate the individual competitiveness and commitment to the medical profession of first year medical students in Erzurum, Turkey.

**Methods:** Students starting the first year of the Faculty of Medicine were asked to answer the questions on the five-point Likert scale between "strongly disagree" and "strongly agree" with the online questionnaire method. The questionnaire included "Medical Faculty Students' Commitment to the Medical Profession Scale" consisting of nine questions and the "Revised Individual Competitiveness Scale" consisting of 11 questions with reversely coded three questions. The data obtained were statistically analyzed using Student t-test, one-way ANOVA and Pearson correlation tests.

**Results:** Out of 517 people who accepted to participate in the study and answered the questionnaire, 151 questionnaires with incomplete, missing or duplicate results were excluded from the study. Of the 366 participants included in the study, 54.1% (n = 198) were female, 45.9% (n = 168) were male; Their average age was  $19.1 \pm 1.3$  years. The average individual competition scale score of the participants was  $33.21 \pm 3.38$ ; The scale of commitment to the medical profession was found to be  $37.16 \pm 5.89$ . While the commitment score of female participants to the profession was  $38.63 \pm 4.97$ , it was found to be  $35.44 \pm 6.41$  for men, and a statistically significant difference was found ( $p < 0.001$ ). While the individual competitiveness scale score of the female and male participants were  $33.17 \pm 3.38$ , and  $33.24 \pm 3.39$ , respectively, no statistically significant difference was found ( $p = 0.839$ ). There was no statistically significant difference between the scores of both scales in terms of age ( $p > 0.05$ ). A weak positive correlation was found between the individual competition scale score of the participants and their commitment to the profession ( $p = 0.030$ ;  $r = 0.114$ ).

**Conclusion:** In this study, for the first time in the literature, the individual competitiveness and commitment to the medical profession of the Medical Faculty students in Turkey were investigated. It was found that there was a positive correlation between the scores of the individual competitiveness and commitment to the medical profession, and female physician candidates were found to be more dedicated to the medicine profession than males.

**Keywords:** *Individual Competitiveness Index; Commitment to Profession of Medicine; Medical students.*

# Evaluation of staging 18F FDG PET/CT images with conventional and volumetric data in patients diagnosed with rectal cancer: Do the characteristics of the primary tumor provide information about the metastatic potential of the tumor?

Dr. Nazlı Pınar Karahan Şen<sup>1</sup> and Prof. Gamze Çapa Kaya<sup>2</sup>

Dokuz Eylul University, Faculty of Medicine Department of Nuclear Medicine, Izmir, Turkey <sup>1,2</sup>  
E-mail <sup>1</sup>: drpinarkarahan@hotmail.com, E-mail <sup>2</sup>: gamze.capa@gmail.com

## Abstract

**Objective:** In 18F FDG PET/CT imaging, in addition to the conventional standard uptake value (SUV) parameters of the tumor tissue, metabolic tumor volume (MTV) and tumor lesion glycolysis (TLG) values can be obtained. In this study, we investigated the relationship between the conventional and volumetric characteristics of the primary tumor obtained from the staging F-18 FDG PET/CT images and the metastatic potential of the primary tumor at the time of diagnosis in patients diagnosed with rectal cancers.

**Method:** Fifty-seven patients who were diagnosed with rectal cancer and underwent 18F FDG PET/CT for staging between January 2012 and January 2021 were included in our study. Primary tumor, regional lymph node metastasis, distant metastasis and KRAS mutation were examined. Conventional and volumetric data of the tumor were obtained by drawing the area of interest (ROI) of the primary tumors from PET/CT images. In order to prevent manual error, the area of interest was drawn in the tumor area with a minimum SUV value of 2 and above. SUV values (minimum, maximum, standard, mean, peak) and metabolic data (MTV and TLG) of the primary tumor within the drawn ROI were obtained. Patients were evaluated in two different groups as KRAS mutant and wild-type. A statistically significant difference in terms of SUV, TLG and MTV values in patients with or without lymph node metastasis in PET/CT and patients with distant metastasis and no metastasis was investigated by Mann Whitney U test.  $P < 0.05$  was considered significant. Receiver Operating Curve (ROC) analysis was performed for the significant features.

**Results:** While 26 of the patients were KRAS mutant, 31 patients were wild-type. Lymph node metastasis was found in 66.7% of the patients, and distant metastasis was found in 61.4%. There was no significant difference in lymph node metastasis and distant metastasis levels between patients with and without KRAS mutation ( $p > 0.05$ ). In KRAS mutant patients, no significant difference was found between the conventional and volumetric characteristics of the primary tumor and the observation of lymph node and distant metastasis. However, TLG and MTV values of the primary tumor were significantly higher in patients with lymph node (AUC: 0.746, 0.737;  $p=0.023$ ,  $p=0.029$ ), and distant metastasis (AUC: 0.718, 0.732;  $p=0.048$ ,  $p=0.035$ ), in patients with KRAS wild-type.

**Conclusion:** Although rates of local and distant metastasis at the time of diagnosis are similar in patients diagnosed with KRAS mutant and wild-type rectal cancer, in this study, it was observed that the high TLG and MTV values of the primary tumor in KRAS wild-type patients were associated with lymph node and distant metastasis at the time of diagnosis. However, in KRAS mutant patients, there was no relationship between conventional and volumetric features and metastasis at the time of diagnosis.

**Keywords:** 18F-FDG PET-CT, MTV, Rectal cancer, KRAS.

# Is the Ganglion Cell Layer Thickness more sensitive marker than retinal nerve fibre layer thickness in Multiple Sclerosis Patients?

Dr. Nuray Can Usta<sup>1</sup> and Dr. Betül Önal Günay<sup>2</sup>

Health Sciences University, Trabzon Training and Research Hospital, Neurology Clinic, Trabzon, Turkey <sup>1</sup>

Health Sciences University, Trabzon Training and Research Hospital, Eye Clinic, Trabzon, Turkey <sup>2</sup>

E-mail <sup>1</sup>: dr.nuraycan@hotmail.com, E-mail <sup>2</sup>: drbetulonal@yahoo.com

## Abstract

**Background:** Optical coherence tomography (OCT) can be used in the follow-up of Multiple Sclerosis (MS), which is the most common cause of non-traumatic disability in young adults.

**Purpose:** It was aimed to evaluate the OCT findings of Multiple Sclerosis patients who applied to the MS outpatient tertiary referral hospital clinic in April 2021 and those in the healthy control group.

**Methodology:** Patients who applied to the MS outpatient clinic of the tertiary referral hospital between April 1, 2021 and April 28, 2021 were included in the study. The exclusion criteria were patients who failed to comply with the OCT examination and patients who did not accept the tests. Age-matched healthy individuals were selected as control group. The patients underwent a comprehensive ophthalmological and neurological examination and OCT parameters were compared.

**Results:** A total of 34, including 12 MS patients and 22 healthy controls, were included in the study. Three of the patients were male and 9 of them were female in MS. The mean age of MS patients was  $36.5 \pm 11.6$  (19-65), and the mean age of the control group was  $35.5 \pm 7.2$  and there was no significant difference in terms of age ( $p=0.79$ ) and gender ( $p=0.8$ ). There was no significant difference between patients who had previously optic neuropathy and had not ( $p<0.05$ ). Compared to the control group, there was no significant difference in Retinal Nerve Fibre Layer (RNFL) thickness on all quadrants in MS patients. Compared to macular thickness (MT), all quadrants except central MT were significantly lower in MS patients [central MT in MS versus control group ( $258.9 \pm 19.6$  versus  $266.9 \pm 15.4$ ;  $p=0.21$ ), superior MT ( $332.3 \pm 17.9$  versus  $350.8 \pm 8$ ;  $p=0.007$ ), temporal MT ( $314.9 \pm 16.2$  versus  $335.1 \pm 11.2$ ;  $p=0.001$ ), inferior MT ( $330.7 \pm 20.3$  versus  $346.7 \pm 11.6$ ;  $p=0.01$ ), nasal MT ( $335.1 \pm 20.7$  versus  $350.7 \pm 13.1$ ;  $p=0.02$ )]. Furthermore in correlation with central MT, thinning was found in the ganglion cell layer (GCL) in all areas except the central region compared to the control group [central GCL ( $13.6 \pm 3.1$  versus  $14.8 \pm 3.0$ ;  $p=0.45$ ), superior GCL ( $48.9 \pm 6.6$  versus  $55.4 \pm 3.3$   $p=0.006$ ), temporal GCL ( $42.4 \pm 7.6$  versus  $42.4 \pm 7.6$ ;  $p=0.008$ ), inferior GCL ( $49.1 \pm 5.5$  versus  $55.6 \pm 3.4$ ;  $p=0.001$ ), nasal GCL ( $48.6 \pm 5.8$  versus  $55.0 \pm 3.9$ ;  $p=0.005$ )]

**Conclusion:** In MS patients, in the ophthalmology department, patients are generally evaluated and followed up with RNFL thickness map. In our study, it was found that there was no significant difference in RNFL thickness map values in MS patients compared to healthy controls, but there was a significant difference in the GCL. It was also thought that the decrease in central MT might be related with thinning in the GCL. Therefore, GCL may be a more sensitive surrogate marker in the follow up of MS patients. Prospective studies involving a greater number of patients are needed to ascertain the relationship between GCL thickness and MS.

**Keywords:** Ganglion Cell Layer, Multiple Sclerosis, Optic Cohorence Tomography, Retinal Nerve Fibre Layer

# How can the ocular surface cells be evaluated histologically with the impression cytology method?

**Dr. İrem İnanç<sup>1</sup>, Dr. Burcu Kazancı<sup>2</sup>, Dr. Fatma Çorak Eroğlu<sup>3</sup> and Prof. Bizden Sabuncuoğlu<sup>4</sup>**

*Ankara University, Faculty of Medicine, Department of Histology and Embryology, Ankara, Turkey<sup>1,4</sup>  
Ulucanlar Eye Research and Education Hospital, Department of Ophthalmology, Ankara, Turkey<sup>2,3</sup>  
E-mail<sup>1</sup>: iinanc@ankara.edu.tr, E-mail<sup>2</sup>: drburcus@hotmail.com, E-mail<sup>3</sup>: dr\_fatoscorak@hotmail.com,  
E-mail<sup>4</sup>: sabuncu@medicine.ankara.edu.tr*

## Abstract

**Aim:** Impression cytology refers to the contact of filters such as cellulose acetate, polycarbonate, nitrocellulose or polyethersulfone to the ocular surface and taking cells from this area. It is possible to obtain superficial tissues using a non-invasive impression cytology technique without a biopsy to evaluate ocular surface diseases. In this study we aimed to evaluate the samples obtained from volunteers with the impression cytology technique in terms of histology.

**Method:** Samples were removed from the conjunctiva by rubbing on a filter paper with a diameter of 5x10mm from 20 volunteers. The samples were fixed in 96% alcohol. The cells on the filter paper were stained with Periodic acid-Schiff (PAS) and Hematoxylin. Goblet cells stained with PAS and epithelial cells stained with Hematoxylin. After fixation staining procedure includes following steps: 1. Washing with distilled water(3-5min.), 2. Administration of Periodic Acid(5min.), 3. Rinse in distilled water , 4. Administration of Schiff reagent(5-15min.), 5. Washing in tap water(5min.), 6. Staining with Hematoxylin(1min.), 7. Washing with distilled water(2-3min.), 8. Transition through 96% and 100% alcohol series (1min.for each) and making transparent in xylol (2-3min.), 9. Taking the filter paper on a slide and covering it with coverslip using entellan.

**Results:** We grouped cells in line with the images we obtained.

According to the epithelial cell appearance:

1. Group: The cells have round shape, the nucleus are large and cover the cytoplasm to a large extent.
2. Group: The cells have polygonal shape and the nucleus are smaller than first group.
3. Group: The cells have polygonal shape and pycnotic nucleus.

According to the Goblet cell appearance:

1. Group: Goblet cells are in round or oval shape. They stained with PAS in large numbers.
2. Group: Goblet cells are in round or oval shape. They are in few numbers.
3. Group: There is no Goblet cells.
4. Group: There is abundant content stained with PAS that is thought to be Goblet cell secretions.
5. Group: There is few content stained with PAS that is thought to be Goblet cell secretions.

**Conclusion:** In this study, when comparing the anamnesis received from patients and examining the ocular surface we determined that various ocular surface irritation findings were found higher in those with polygonal shape of epithelial cells, smaller or pycnotic nuclei and low numbers of goblet cells. publication since 2016. This increase in research has built a worldwide network of scientists participating in intra-national and international collaboration.

**Keywords:** *impression cytology, ocular surface, conjunctiva epithelium, Goblet cell*

# Does the tissue morphology change during the development of the age-related metabolic syndrome?

**Mihriban Alemdar, M.Sc<sup>1</sup>, Dr. İrem İnanç<sup>2</sup>, Assoc. Prof. Deniz Billur<sup>3</sup> and Prof. Dr. Nuray Yazıhan<sup>4</sup>**

*Ankara University, Institute of Health Sciences, Interdisciplinary Food, Metabolism and Clinical Nutrition Department, Ankara, Turkey<sup>1</sup>*

*Ankara University, Faculty of Medicine, Department of Histology and Embryology, Ankara, Turkey<sup>2,3</sup>*

*Ankara University, Faculty of Medicine, Internal Medicine, Department of Pathophysiology, Ankara, Turkey<sup>4</sup>*

*E-mail<sup>1</sup>: mihribanalemdar@hotmail.com, E-mail<sup>2</sup>: iinanc@ankara.edu.tr, E-mail<sup>3</sup>: denizbillur@gmail.com, E-mail<sup>4</sup>: nurayyazihan@yahoo.com*

## Abstract

**Aim:** Studies show that the prevalence of metabolic syndrome (MetS), which is progress with abdominal obesity, insulin resistance, glucose intolerance, hypertension and dyslipidemia increases in the elderly population. Changes are occurred in the immune system with aging, deficiencies in the immunological response to aging-related pathogens are observed. Elderly population in the world and Turkey and the prevalence of diseases in the elderly population increased, it is thought that the effect of metabolic syndrome development process of developing immune deficiency with aging. Based on this, we aimed to evaluate the liver and kidney tissues morphologically in rats with adult, aged and age-related metabolic syndrome in this study.

**Method:** In this study, we used adult age (6-month) and aged (24-month) Wistar-type male rats. Oral Glucose Tolerance Test used to evaluate insulin resistance and blood glucose levels above 100 mg/dl were accepted as the metabolic syndrome group. In this context; adult, aged MetS(-) and aged MetS(+) groups were formed in our study. At the end of the experiment, the rats liver and kidney tissues were removed under anesthesia and fixed in a 10% phosphate buffered formalin solution for 72 hours. Paraffin blocks were obtained by routine histological tissue processing method and then 5 µm thick sections were taken with Leica RM 2125RT model sliding microtome. Hematoxylin-Eosin and Mallory Azan (MA) stainings were applied to the sections.

**Results:** In adult group, classical liver lobules were observed in the usual appearance. In aged MetS (-) and aged MetS (+) groups, collagen fibers increase was observed around the vena centralis and sinusoid structures and the sinusoids were also enlarged. In the portal area; enlargement of bile ducts, an increase in the number of hepatocytes containing pycnotic nuclei, congestion and local mononuclear cell infiltration were observed especially in the aged MetS (+) group. When the aged MetS (-) and aged MetS (+) groups were compared with the adult group, shrinkage and fragmentations were observed in the kidney glomerular structure and expansion in the Bowman's space was also determined. The increase in the collagen fibers with MA staining in the glomerular capillaries was evaluated as thickening in the glomerular basement membrane. In the same sections, collagen increase was observed in the parietal layer of the Bowman capsule. In addition, an increase in inflammation was observed in the aged MetS (+) group.

**Conclusion:** In aged groups; liver and kidney tissue morphology differed from the adult groups. Edematous areas, congestion, mononuclear cellular infiltration, and collagen fibers were significantly increased in both aged groups. Especially the cell inflammation was distinctive in the aged MetS (+) groups. For this reason, we aimed to evaluate the phenomenon of inflammation in terms of macrophage polarization in our future studies.

**Keywords:** *Aging, Metabolic Syndrome, Liver, Kidney, Morphology.*

## Histological evaluation of the effectiveness of different microorganism types forming infections in the abscess formation

**Dr. Mehmet Batu Ertan<sup>1</sup>, Dr. İrem İnanç<sup>2</sup>, Dr. Mehmet Yağız Ayduğan<sup>3</sup>,  
Assoc. Prof. Ebru Evren<sup>4</sup>, Prof. Esra Erdemli<sup>5</sup> and Prof. Bülent Erdemli<sup>6</sup>**

Ankara University, Faculty of Medicine, Department of Orthopedics and Traumatology, Ankara, Turkey<sup>1,3,6</sup>  
Ankara University, Faculty of Medicine, Department of Histology and Embryology, Ankara, Turkey<sup>2,5</sup>  
Ankara University, Faculty of Medicine, Department of Medical Microbiology, Ankara, Turkey<sup>4</sup>  
E-mail<sup>1</sup>: mbatuertan@gmail.com, E-mail<sup>2</sup>: iinanc@ankara.edu.tr, E-mail<sup>3</sup>: aydugan@ankara.edu.tr, E-mail<sup>4</sup>:  
evren@ankara.edu.tr, E-mail<sup>5</sup>: erdemli@medicine.ankara.edu.tr, E-mail<sup>6</sup>: berdemli@medicine.ankara.edu.tr

### Abstract

**Aim:** Total knee and hip arthroplasties are safe and effective surgical procedures. It decreases the patients' complaints, restores movement functions and improves quality of life, especially in the elderly population. The risky side of this process is that microorganisms around the prosthesis accumulate and create a biofilm over time and in the end they can cause infection. Despite the increased interest and developments in the diagnosis and treatment of prosthetic joint infection, this topic is still a mystery. In this study, we designed a method similar surgical procedure mentioned above on a rat model and we aimed to determine if there is any difference between the types of microorganisms in biofilm formation histologically.

**Method:** 6 standard titanium alloy (Ti6Al4V) discs were placed in pouch regions formed in the interscapular regions of the experimental animals and each pouch were infected with microorganisms (except group 1) such as staphylococcus epidermidis (group 2), staphylococcus aureus (group 3), pseudomonas aeruginosa (group 4) and candida albicans (group 5). The concentration of the each microorganism was 1ml of  $1.5 \times 10^7$  cfu/ml. In the 2nd, 4th and 6th weeks of the study, 2 discs were removed from each group of animals and examined in the scanning electron microscope. When the experimental procedure was completed at the 6th week, the rats were sacrificed, the abscess tissues were taken in 10% buffered formaline, and paraffin blocks obtained by the routine histological processing method. Sections in 4µm thickness were taken and the cyst area was evaluated with Hematoxylin&Eosin staining. The spread and involvement of these microorganisms on titanium discs in in vitro conditions were investigated with scanning electron microscope.

**Results:** When compared with the control group, the microorganisms development in in vitro conditions on titanium discs determined with the scanning electron microscopy. Except the control group intensive inflammation seen in all groups. Consequently, there was excessive amount of macrophages and leukocyte serial cells. Inflammatory cell infiltration, vessel formation and edema areas that indicates inflammation was determined with Hematoxylin&Eosin staining. Furthermore, significant connective tissue increasing was observed.

**Conclusion:** We determined intense inflammation in all groups where microorganisms applied with this experimental model and we also detected effective defence cells against inflammation. Although there was no difference between the groups, the tissue regeneration has better morphology with Hematoxylin&Eosin staining in staphylococcus aureus treated groups. Therefore, the molecular mechanisms underlying the stages of inflammation and tissue regeneration is planned with our future studies.

**Keywords:** Arthroplasty surgery, infection, microorganisms, inflammation, histology

# Ultrastructural examination of anterior capsule samples taken from patients with congenital or juvenile cataract cases

**Dr. İrem İnanç<sup>1</sup>, Dr. Pınar Bingöl Kızıltunç<sup>2</sup>, Dr. Ferhad Özer<sup>3</sup>,  
Prof. Huban Atilla<sup>4</sup>, and Prof. Belgin Can<sup>5</sup>**

*Ankara University, Faculty of Medicine, Department of Histology and Embryology, Ankara, Turkey <sup>1,5</sup>*

*Ankara University, Faculty of Medicine, Department of Ophthalmology, Ankara, Turkey <sup>2,3,4</sup>*

*E-mail <sup>1</sup>: iinanc@ankara.edu.tr, E-mail <sup>2</sup>: pinarbingol84@gmail.com, E-mail <sup>3</sup>: ferhad\_ozer@hotmail.com, E-mail <sup>4</sup>:*

*hatilla@medicine.ankara.edu.tr, E-mail <sup>5</sup>: belgincan@yahoo.com*

## Abstract

**Aim:** Cataract is the loss of transparency of the lens and it is mainly a pathology in elder population. However, it can be seen in congenital and juvenile forms. In this study we aimed to examine the histopathological changes occurred in the anterior lens capsule samples from congenital or juvenile cataract patients.

**Method:** Tissue samples taken from 10 patients were fixed in 2.5% glutaraldehyde with 0.1 M phosphate buffer at +4 °C for 2-4 hours. Samples washed in 0.1 M phosphate buffer and there after treated with osmium tetroxide for 2 hours. Block staining was performed with uranyl acetate. For dehydration, samples were passed through the alcohol series and were embedded in Araldite 6005 material. Semi-thin sections in 800 nm thickness taken in Leica ultracut R ultramicrotome were stained with toluidine blue-azur II and sections in 60 nm thickness were stained with uranyl acetate and lead citrate. Sections examined and viewed with LEO 906 E transmission electron microscope.

**Results:** At the electron microscopic level, vacuolized areas appeared between capsule and subcapsular epithelium, whereas in some of the samples the lens capsule and the subcapsular epithelium were observed in normal formation. The regular subcapsular epithelium is composed of a single layer of cuboidal cells with round nuclei. In our samples accompanying to the changes of the cellular shapes, different types of nuclei such as flat, oval forms were observed in the subcapsular epithelium. In some of the cases the normal structure of the single-layer epithelium was replaced by the multi-layer epithelial form. The cytoplasmic organelles were degenerated in most of the epithelial cells and the most prominent damage was determined in mitochondria which showed abnormal structure. They were larger in size and loss of cristae was evident. Within all samples only one case had connective tissue instead of epithelial cells.

**Conclusion:** In this study, vacuolized areas were observed in the lens anterior capsule tissue of both congenital and juvenile cataracts. Especially different forms of epithelium and degeneration of organelles were significant in juvenile cataract. We observed these changes in all cataract cases. On case basis, the ultrastructural changes were not specific to define the diseases with the electron microscope. The number of patients need to be increased to get significant results.

**Keywords:** *Congenital and juvenile cataract, anterior lens capsule, electron microscopy.*



# The Psychosocial Effects of The Covid-19 Pandemic on Adolescents at A Private High School in Istanbul

**Dr. Gokce Hazar Otcu<sup>1</sup>, Aydin Arman Canbaz<sup>2</sup>, Sarp Esen<sup>3</sup>, and Assoc. Prof. Meryem Merve Oren<sup>4</sup>**

*Istanbul University Istanbul Faculty of Medicine, Department of Public Health, Istanbul, Turkey <sup>1,4</sup>*

*High school students, GIB program, Istanbul, Turkey, <sup>2,3</sup>*

*E-mail <sup>1</sup>: gh.otcu@istanbul.edu.tr, E-mail <sup>2</sup>: armancnbz1@gmail.com, E-mail <sup>3</sup>: sarpesen@yahoo.com,*

*E-mail <sup>4</sup>: merve.oren@istanbul.edu.tr*

## Abstract

Many countries have closed schools in order to prevent the spread of COVID-19. As a result, adolescents have become removed from the peer and teacher interactions of school life and/or become socially isolated. The aim of this cross-sectional study was to determine the psychosocial effects of the COVID-19 pandemic on students attending a private high school in Istanbul in the 2020-2021 academic year.

Between November 1, 2020 and February 1, 2021, participants completed an online questionnaire including the COVID-19 Phobia Scale (C19P-S), Trait Anxiety Inventory (TAI), and questions about their sociodemographic characteristics, COVID-19 prevention measures taken, and pandemic-related life changes. The minimum sample number was calculated as 199 students, and 92% (n=183) of this target was achieved.

Of the participants, 144 (78.7%) stated that they had very good knowledge of how to prevent the spread of COVID-19, 181 (98.9%) reported wearing face masks for protection, 159 (86.9%) washed their hands for at least 20 seconds, and 140 (76.5%) followed social distancing rules. The most common changes in their lives during the pandemic were increased use of digital devices, feeling socially isolated, and lethargy and laziness due to staying at home for a prolonged period. Based on the group medians, 39 (21.3%) of all participants were found to have COVID-19 phobia, with phobia in the psychological, somatic, social, and economic domains in 74 (40.4%), 57 (31.1%), 66 (36.1%), and 47 (25.7%) of participants, respectively. Based on the median TAI score, 55 (30.1%) of the participants had anxiety.

Adolescents are a vulnerable group; to protect their mental health during the pandemic, risks must be reduced and preventive mental health interventions increased. Parents should provide guidance to adolescents in order to provide access to sources of accurate information, use the internet to support individual and social development, and create alternative ways for peer interaction to reduce isolation and loneliness.

**Keywords:** *Adolescent; COVID-19*

# Evaluation of the effect of mesenchymal stem cell administration on prognosis in critical COVID-19 patients in the intensive care unit

**Dr. Seyfi Kartal**

*Health Sciences University, Trabzon Kanuni Training and Research Hospital, Anesthesiology and Reanimation Clinic, Trabzon, Turkey*

*Email: drseyfikartal@gmail.com*

## Abstract

**Background:** In the literature, there are several publications regarding the use of Mesenchymal stem cells for acute respiratory distress syndrome (ARDS) related to COVID, there is no large-scale research about its use in critical cases.

**Purpose:** In this study, it was aimed to evaluate the effectiveness of MSC application in critical COVID-19 patients who were treated in intensive care unit (ICU)

**Methodology:** All patients included in the study were routinely treated according to the COVID-19 treatment algorithm of the Turkish Ministry of Health; (1) oxygen supplements, (2) antiviral agents, (3) antibiotic agents, (4) glucocorticoid agents.

After 7-10 days of routine treatment patients were included; increase in oxygen demand while under treatment and rapid progression on radiological imaging.

MSC for patients was calculated as a single dose,  $1 \times 10^6$  cells per kilogram as well as with routine treatment. MSC was infused intravenously in 100 ml normal saline for 40 min. Demographic data and various data of the patients were evaluated.

**Results:** In this study, we evaluated 81 ICU patients. There were 49 patients in the mortal group and 32 patients in the survival group in the study. The mean age of mortal group was  $61.35 \pm 12.59$ , and the mean age of the survival group was  $51.19 \pm 12.62$  and there was no significant difference between the age groups ( $p > 0.05$ ). There was no significant difference between the age groups in terms of Body Mass Index, Acute Physiology and Chronic Health Evaluation-II and Modified Charlson Comorbidity Index ( $p > 0.05$ ). Mortality was higher in patients who received invasive mechanical ventilator therapy at the time of MSC application ( $p < 0,001$ ).

Compared to MSC treatment day, partial pressure of arterial oxygen: percentage of inspired oxygen (PaO<sub>2</sub>/FiO<sub>2</sub>) ratio was higher in the survival group ( $59,52 \pm 23,32$  and respectively  $80; 47 \pm 29,62, 92,47 \pm 27,12, 115,33; \pm 34,1$ ) compared to mortal group ( $69,11 \pm 22,79$  and respectively  $78,53 \pm 29,3, 88,07 \pm 27,3, 93,02 \pm 33,8$ ) at days 3, 7, 14 days ( $p < 0,005$ ). There was no difference between the groups in terms of total days of stay in ICU, total days of hospitalization and days stay in ICU after application ( $p > 0,05$ ).

**Conclusion:** ARDS develops in COVID patients due to uncontrolled inflammation and alveolar membrane damage due to cytokine storm. MSC has anti-inflammatory, immunomodulatory, anti-fibrotic, high proliferation, regeneration and tissue repair effects, as well as cleaning pathogens and reducing the damage they cause. After a single dose injection of MSC, pulmonary function and patients' symptoms (PaO<sub>2</sub> / FiO<sub>2</sub> ratio) improved. This increase is not significant enough in all patients. We are of the opinion that MSC treatment has a positive effect on the early stage of the cytokine storm, before the initiation of invasive mechanical ventilation treatment, and reduces the mortality. Further detail research is needed on the subject.

**Keywords:** COVID-19, stem cell, intensive care unit

# Polymicrobial Sepsis Causes Edema and Neuronal Damage in Rat Cerebral Cortex; an Experimental Study

Asst. Prof. Songül Doğanay<sup>1</sup> and Asst. Prof. Özcan Budak<sup>2</sup>

*Sakarya University, Department of Physiology, Faculty of Medicine, Sakarya, Turkey*<sup>1</sup>

*Sakarya University, Department of Histology and Embryology, Faculty of Medicine, Sakarya, Turkey*<sup>2</sup>

*E-mail*<sup>1</sup>: songuldoganay@sakarya.edu.tr, *E-mail*<sup>2</sup>: ozcanbuda@sakarya.edu.tr

## Abstract

**Objective:** Inflammation is thought to play an important role in the pathogenesis of many neurological diseases in recent years. Neuroinflammation is a common finding in acute neurological events such as stroke, trauma and infection, as well as chronic neurodegenerative diseases such as Alzheimer's, Parkinson's, and multiple sclerosis. Sepsis defined as an inflammatory response, and it involves organ systems remote from the locus of the initial infectious insult. Perforation in the intestinal barrier plays an important role in multiple organ failure due to sepsis. In line with this information, the aim of our study was to investigate the damage to the brain tissue due to polymicrobial sepsis.

**Materials and Methods:** Experimental animals (Wistar-albino rats; n = 14) were divided into 2 groups as sham (control) and sepsis group (CLP). Sepsis was induced through the cecal ligation and perforation method (CLP). Twenty-two hours after surgery, brain tissues were taken for histological evaluation. The sections taken from paraffin blocks of brain tissues were stained with hematoxylin and eosin (H&E) and examined histopathologically under a light microscope.

**Results:** In the histomorphological evaluation performed in the experimental groups, serious damage was observed in the cerebral cortex of septic rats. We observed that the cerebral cortex areas of the rats in the control group had regular morphology. However, in CLP group rats, most of the neurons were shrunken and scattered in the cerebrum. Severe degeneration, perineuronal and perivascular edema and inflammatory cell infiltration were observed in neurons with pyknotic nuclei. These changes were found to be statistically significant when compared to the control group (p <0.001).

**Conclusion:** According to the results of our study, polymicrobial sepsis causes serious damage to the cerebral cortex.

**Keywords:** Brain injury, cerebral cortex, neuronal degeneration, sepsis.

# Pediatric Endocrinology and Syrian Refugee Children: A Photo We Want to Look Together

**Dr. Tuğba Kontbay<sup>1</sup> and Dr. Müge Atar<sup>2</sup>**

*Şanlıurfa Training and Research Hospital<sup>1,2</sup>*

*E-mail<sup>1</sup>: tugbakontbay@gmail.com , E-mail<sup>2</sup>: drmugeatar@gmail.com*

## **Abstract**

**Aim:** Increasing number of refugees take concern of health workers, and lead to identify health challenges in refugee population. Along with the follow-up characteristics of the children under the current life conditions, the evaluation of the newly diagnosed children in our country is important for the future health goals and plans.

In this study, we aimed to determine the etiological distribution of refugee children who applied to our outpatient clinic.

**Material-Methods:** Refugee children admitted to Pediatric Endocrinology outpatient clinic during July-December 2019 was evaluated in this cross-sectional study. Hospitalized patient were excluded. The cases were grouped according to their age, gender, anthropometric measurements, reasons for admission, diagnosis whether they had a chronic disease at the time of admission, and the patients were photographed in terms of endocrine diseases.

**Results:** The mean age of the 414 patients were  $8.13 \pm 5.1$  years. 62.5% of patients were applied due to short stature while remaining complaints were associated with thyroid disease (20.1%), diabetes mellitus (5.3%), pubertal disorders (2.7%), obesity (2.7%), rickets (2.4%), and the others (4.4%). After pediatric endocrinology evaluation, the patients diagnosed with short stature (33.9%), malnutrition (17%), congenital hypothyroidism (6.1%), diabetes mellitus (5.1%), subclinical hypothyroidism (3.4%), obesity (3.1%), calciopenic rickets (2.4%), Hashimoto thyroiditis (2.2%), precocious puberty (1.2%), others (6.5%) while 17.7% of the patients were healthy.

**Conclusion:** The evaluation of refugee children who have been diagnosed in their country or newly diagnosed in our country is very important for the world health goals. This study is designed to raise awareness on a serious condition.

**Keywords:** *Refugee Children; Pediatric Endocrinology; etiological distribution*

# Methanol Intoxication with Cerebral Hemorrhage: A Case Study

**Dr. Tuğçehan Sezer Akman<sup>1</sup>, Dr. Hale Kefeli Çelik<sup>2</sup> and  
Assoc. Prof. Dr. Ahmet Şen<sup>3</sup>**

*Health Sciences University, Anesthesiology and Reanimation, Samsun Training and Research Hospital, Samsun, Turkey<sup>1,2,3</sup>  
E-mail<sup>1</sup>: tgchnszr@gmail.com, E-mail<sup>2</sup>: ck\_hale@gmail.com, E-mail<sup>3</sup>: ahmetsenau@gmail.com*

## Abstract

Methyl alcohol (methanol, wood alcohol) is the simplest alcohol used in manufacture of various laboratory and industrial products. Methanol intoxication is a serious case potentially ending up with mortality in case of late diagnosis and treatment. In present case, 65 years old male patient was brought to emergency service with the complaints of confusion and epigastric pain. The patient with a learned cologne drinking history had deep metabolic acidosis. The patient's history and current metabolic table bring into mind the methanol intoxication. Intravenous sodium bicarbonate, ethanol and 5% dextrose were administered. Cardiopulmonary arrest was developed, resuscitation was applied to patient and then hemodialysis was applied accompanied with inotropic support. Following the dialysis, the patient was treated in an intensive care unit in intubated and sedated fashion. Following the termination of sedation of the patient with improved metabolic table, Glasgow coma score (GCS) remained low and tracheotomy was opened and follow ups continued under ventilator support. In brain BT taken in 15th day of hospitalization, bilateral intracerebral hemorrhage areas were seen and the patient was lost in 20th day of hospitalization. Accompanied with the present case, the objective of the study was to evaluate diagnosis, treatment and follow up of methanol intoxication.

**Keywords:** *Methyl Alcohol; Poisoning; Metabolic Acidosis; Intoxication.*

# Is There a Relationship between Hip Fracture Surgical Treatment and Developing Dementia? One Year- Prospective Study

**Dr. Tuğrul Ergün**

*Ministry of Health Mardin State Hospital, Mardin, Turkey  
E-mail: drergun14@gmail.com*

## Abstract

**Background:** Hip fractures are common pathologies with high mortality and morbidity in the elderly people (1). Its treatment can be performed with different surgical options for early weight bearing and mobilization (2). Many studies have shown that patients with hip fractures are susceptible to developing dementia in their long-term follow-up (3).

**Purpose:** In this study, we aimed to investigate the relationship between two different surgical techniques which are commonly applied in the treatment of hip fractures and the development of dementia.

**Methodology:** Patients who underwent orthopedic surgery for hip fractures were included in the study. Patients with advanced dementia and who could not be followed up postoperatively period were excluded from the study. The demographic characteristics of the patients (age, gender), the surgical technique [Hemiarthroplasty and Proximal Femoral Nail (PFN)] and the Mini-Mental State Examination (MMSE) test that was used to evaluate the patients for dementia during their stay in the hospital and 12 months after the operation were recorded. The Mann-Whitney U test was used to analyze the relationship between two different surgical techniques and patient's dementia.

**Results:** 31 patients were included in the study. The mean follow-up time period was 15 (13-23) months. While 61.3% (n = 19) of the patients were female, the mean age was  $77.1 \pm 9$  years. 54.8% (n = 17) of the patients were treated with PFN. Firstly, the mean MMSE value was  $17.7 \pm 1$  in the hemiarthroplasty group and it was  $17.8 \pm 1$  in the PFN group. After 1 year, while the mean MMSE value in the PFN group was  $16.4 \pm 1$ , it was  $13.7 \pm 1$  in the hemiarthroplasty group. While a decrease was observed in the mean MMSE values of both groups, this decrease was observed more than in the hemiarthroplasty group compared to the PFN group. ( $p < 0.5$ ).

**Conclusion:** The patient group who underwent hemiarthroplasty in the treatment of hip fractures should be followed closely for the development of dementia, and minimally invasive surgical techniques may be considered in the hip fracture treatment.

**Keywords:** Hip Fracture, Dementia, Hemiarthroplasty, Proximal Femoral Nail (PFN).

# Effects of Virtual Reality on Stress Management of Disasters

Volkan Ülker, MD, PhD<sup>1</sup>, Assoc. Prof. Bedia Gülen<sup>2</sup> and Prof. Yusuf Yürümez<sup>3</sup>

*Sakarya University Training and Research Hospital, Clinics of Emergency Services, Sakarya, Turkey<sup>1</sup>*  
*Bezmialem Foundation University, Institutes of Health Sciences, Department of Disaster Medicine, Istanbul, Turkey<sup>1</sup>*  
*Medipol University Medical Faculty, Department of Emergency Medicine, Istanbul, Turkey<sup>2</sup>*  
*Sakarya University Medical Faculty, Department of Emergency Medicine, Sakarya, Turkey<sup>3</sup>*  
E-mail<sup>1</sup>: volkanulker07@yahoo.com, E-mail<sup>2</sup>: drbediagulen@yahoo.com, E-mail<sup>3</sup>: yusufyurumez@yahoo.com

## Abstract

**Introduction:** Technology-supported applications used in healthcare services are effective in increasing service efficiency and improving quality, providing fewer errors, while also allowing patients to recover in a shorter time. Virtual reality (VR) applications one of these technologies. VR presents a new human-computer interaction paradigm where users are not external observers on a computer screen but are active participants in a computer-generated three-dimensional virtual world. In this study, it is aimed to use VR in medical triage and to evaluate whether it will provide benefits in terms of stress management for the Healthcare Personnel (HCP) that performs at the scene in disasters, and this report covers the first data obtained from the study.

**Materials and Method:** This study was aimed to be carried out between 01.01.2020-01.01.2021, with a total of 135 participants within Disaster Medicine Department of Bezmialem University. Before the study, permission was obtained from the local ethics committee (Date: 10.09.2019, Decision No: 16/327). The population of the study was orientated from doctors and nurses of Emergency Clinics of Sakarya University Training and Research Hospital and senior students of Sakarya University Faculty of Medicine. Theoretical and video-supported practical lessons about field management, stress management, triage, life-saving interventions and the structure of field hospitals in disasters were given through participants and the stress levels of the participants were measured with Burnout Measure Short Version (BMSV), Perceived Stress Scale (PSS) and Beck Anxiety Inventory (BAI). Then, VR was made about a disaster caused by chemical explosions and participants were asked to triage for the victims. After VR, the stress levels of the participants were again determined with BAI. The data obtained were uploaded to the IBM STATISTICS SPSS computer system and analyzed. As a result of the analysis, values below 0.05 were accepted as significant.

**Result:** According to the first findings obtained from the research, 52 of the participants (n: 80) were women and their mean age was 23.78. The most prominent findings of the participants according to the BMSV are; he was exhausted, insecure and depressed. According to PSS; routine daily difficulties, personal problems, and the desire to succeed are the most prominent findings. While the proportion of participants who had no anxiety in terms of BAI before the training was 30% and the rate of severe anxiety was 12.5%. The proportion of no anxiety after the application was 30%, while severe anxiety rate was 16.3%.

**Conclusion:** We believe that VR practices for HCP responding to disasters did not cause any statistical change in the assessment of stress management at the time of the incident with BAI, but this result may be due to the initial limited number of participants.

**Keywords:** *Disaster, Stress, Triage, Virtual Reality*

## Two siblings with Hyper IgE Syndrome diagnosed by Exome Sequence; DOCK8 mutation

**Asist. Prof. Yeşim Özdemir**

*Uskudar University Medical Faculty, Medical Genetics Department, Istanbul Turkey*

*Memorial Sisli Hospital, Genetic Diagnosis Center, Istanbul, Turkey*

*E-mail: sevdayesim.ozdemir@uskudar.edu.tr, sevdayesimozdemir@gmail.com*

### Abstract

Hyper IgE Syndrome is a syndrome characterized by primary immunodeficiency, eosinophilia, recurrent Staphylococcal infections, recurrent lung infections, eczematous dermatitis, dental and skeletal system abnormalities. While the autosomal dominant form occurs with heterozygous mutations in the STAT3 gene, the autosomal recessive form is caused by homozygous or compound heterozygous mutations in the DOCK8 gene. Autosomal recessive form is associated with severe eczematous and viral infections and can lead to skin cancers and premature death.

6- and 8-years old brothers presented to our clinic with complaints of severe immunodeficiency, eczematous lesions, growth and developmental retardation. Physical examinations showed severe thickening and roughening of the skin, diffuse dental caries and loss of enamel, candidiasis and aphthous lesions in the oral mucosa, perianal abscess, especially in the scrotum skin, in the big one. There was diffuse infiltration and consolidation in the lung and pleural effusion. One of the siblings had minor fatty liver and enlargement. Leukocytosis and eosinophilia were dominant in their peripheral blood smear. In the family history, it was learned that the mother and father were third degree relatives. Dual Exome Sequence analysis performed for etiology and deletion in the DOCK8 gene covering the 46th exon was detected in both siblings. This deletion was confirmed by MLPA method. While the children were followed up in the Pediatric Hematology-Oncology Clinic, preimplantation Genetic Diagnosis was recommended to the family for their next childs. Children with this mutation, which was previously published only once in the literature, were also presented as rare cases.

**Keywords:** *Hiper IgE, DOCK8, immune deficiency, whole exome sequencing, WES*



# Evaluation of protective effects of Ghrelin on Gastric tissue of ovariectomized rats

**Dr. Özlem Tuğçe Çilingir-Kaya**

*Marmara University, School of Medicine, Histology and Embryology Department, İstanbul, Turkey  
E-mail: tugce.cilingir@marmara.edu.tr*

## Abstract

**Background:** Menopause is a significant event that effects women body physiologically. Whereas some pathophysiological effects occur before menopause, some of them occur after menopause named as postmenopausal syndrome. Many studies showed that the gastric inflammation is increased after menopause. Ovariectomy is one of the most commonly used animal models to mimic the human menopause.

**Purpose:** In the present study we aimed to evaluate possible protective effects of ghrelin on gastric tissue in ovariectomized (OVT) rat at light and electron microscopic levels.

**Methodology:** Under ketamine-chlorpromazine anesthesia (100 mg/kg, 0.75 mg/kg), Sprague-Dawley rats (n=12) underwent bilateral OVT, while control group had sham-surgery (n=6). Four weeks after surgery, half of OVT rats were treated intraperitoneally with ghrelin (1 mg/kg/week) for 4 weeks, while others were not treated. Rats were euthanized by cardiac puncture at the end of 8th weeks, and serum levels of estradiol were measured. Gastric tissue was also analyzed with light and scanning electron microscopy. For this purposes, part of tissues were fixed in formaldehyde and underwent routine histological assessments for light microscopic examinations whereas the remaining part of tissues were fixed in glutaraldehyde and coated with gold for scanning electron microscopic investigations. Then all tissues from the different experimental groups were evaluated in terms of histological alterations.

**Results:** Increased body weights in OVT rats ( $p<0.001$ ) recorded at the end of 2 months was not changed with ghrelin. Serum estradiol levels were reduced ( $p<0.05$ ) verifying altered gonadal hormone status. Apical mucus cells and gastric pits were observed in regular morphology in sham group while vascular dilatation and degeneration was observed in OVT rats, whereas in the ghrelin group it was generally observed similar to sham group.

**Conclusion:** In conclusion, mild gastric inflammation in early post-menopausal period appears to be attenuated by ghrelin treatment, and it is required further investigations.

**Keywords:** *Ghrelin, postmenopausal, gastric Inflammation, light microscopy, electron microscopy.*

# Effect of Chronic Alcohol Abuse on the Spectroscopically Determined Blood Secondary Structure of Proteins and Lipid Balance

Zozan Güleken, PhD

Üsküdar University, Department of Physiology, Medical Faculty, Istanbul, Turkey

E-mail: zozan.guleken@uskudar.edu.tr

ORCID: <https://orcid.org/0000-0002-4136-4447>

## Abstract

Alcohol abuse is the most common and costly form of drug abuse. The disease is a major contributing factor leads to many disease categories. The main purpose of this study is to identify the effect of chronic alcohol consumptions in the secondary structure of proteins and lipid balance in the blood samples of the patients. To identify the changes on blood parameters, we measured lipid peroxidation and oxidative load of the alcohol abuse patients (n=20) and healthy subjects (n=20). We measured malondialdehyde (MDA), glutathione (GSH), superoxide dismutase (SOD) and catalase (CAT) level to evaluate oxidative load. Additionally, to see structural changes, we used Fourier InfraRed Spectroscopy (FTIR) tool in the regions of 2nd derivative data between 1250 and 1300 cm<sup>-1</sup>, 1500-1700 cm<sup>-1</sup> and between 1600-1700 cm<sup>-1</sup>. Our results showed that there was an elevated oxidative load with high MDA level (72.13±4.01) compared to the control (25.17±6.13) group (p<0.01). GSH (41.12±7.03) level were elevated compared to the control (7.01 ±3.06) group (p<0.05) too. SOD level was 672.11±56.7 for alcohol abuse patients and 1509.18 ± 78.02 for healthy control (p<0.001). Finally, CAT level was 896.01±40.38 with mean± SD for alcohol abuse patients and 1357±188.96 for healthy control group (p<0.05). FTIR analysis showed that there is an important structural change in the regions of the peaks corresponding to functional groups building nucleic acid, carbohydrates, proteins and lipids structures. Furthermore, asymmetric vibrations of PO<sub>2</sub> groups from nucleic acids (1234 cm<sup>-1</sup>) and CH<sub>2</sub> as well as C=O stretching vibrations of COO<sup>-</sup> groups (1401 cm<sup>-1</sup>) and scissoring vibrations of CH<sub>2</sub> groups from carbohydrates and proteins were observed in FTIR spectra of blood samples. Amide II and amide I, vibrations were placed at 1600-1700 cm<sup>-1</sup> regions, while vibrations of lipids functional groups were noticed respectively. When we compared amide I region obtained for control and alcohol abuse groups, different structure of 1600 cm<sup>-1</sup> – 1700 cm<sup>-1</sup> range, were observed. Consequently, different number of fitted curves, was noticed. These curves were originated from secondary structure of proteins: α-helix and β-sheet. Our results showed that contrarily to laboratory report's Chronic consumption of alcohol causes oxidative load and changes on the structure of blood proteins. Spectroscopic analysis and biochemical analysis are important tools for the diagnosis of the pathophysiologic process of the disease in the level of the changes on biomacromolecules and lipids.

**Keywords:** Alcohol abuse; Physiopathology; Oxidative status; Diagnostic techniques

# Effects of Rosmarinic Acid Against Nephrotoxicity Induced by Cyclophosphamide in Rats

Dilan Çetinavcı, MD<sup>1</sup>, Prof. Engin Yenilmez<sup>2</sup>, Ayşe Firuze Bıyık, MD<sup>3</sup>,  
Prof. Ahmet Alver<sup>4</sup>, Neslihan Sağlam, R.A<sup>5</sup>

*Mugla Training and Research Hospital, Mugla, Turkey <sup>1</sup>*

*Karadeniz Technical University Faculty of Medicine, Histology and Embryology, Trabzon, Turkey <sup>2,3</sup>*

*Karadeniz Technical University Faculty of Medicine, Medical Biochemistry, Trabzon, Turkey <sup>4,5</sup>*

*E-mail <sup>1</sup>: drdilancetinavci@hotmail.com*

## Abstract

In this study, the protective effect of rosmarinic acid (RA) against renal damage caused by cyclophosphamide (CP) was investigated. For this purpose, histopathological and biochemical evaluations were made.

18 Sprague Dawley male rats were randomly divided into three groups: Sham, CP and CP+RA. Sham group was given physiologic serum (PS) by gavage for 8 days from the 1st day of the experiment and PS was administered intraperitoneally on the 8th day of the experiment. CP group was given PS by gavage from the 1st day to the 8th day and a single dose of 200 mg/kg CP was administered on the 8th day of the experiment intraperitoneally. The CP+RA group was given 100 mg/kg RA by gavage for 8 days from first day and 200 mg/kg CP intraperitoneally on the 8th day. On the 9th day of the experiment, the rats were sacrificed and the blood of the rats was taken into the blood tubes. Half of each left kidney removed was taken into a 10% formalin solution for fixation and histological evaluations. Kidney tissue samples were stained using hematoxylin and eosin (H&E) and periodic acid-Schiff (PAS) staining methods for light microscopic evaluation. Tissue MDA, SOD, CAT assessments were made from the other half of the left kidney. BUN and creatinine in blood serum samples were measured.

In histopathological evaluations, tubular atrophy, glomerular damage, vascular congestion, interstitial inflammation, epithelial vacuolization, desquamation in tubular epithelium, leukocyte infiltration and hyaline material accumulation was observed in the kidney sections of the CP group. These disorders were significantly reduced in the CP+RA group compared to the CP group. In PAS stained sections, histopathological findings such as widespread shedding, vacuolization were detected in the tubular epithelium and disruption of brush border and basal membrane were observed in CP group. These disorders decreased in CP+RA group compared to CP group. Increased BUN value in CP group decreased in CP+RA group; whereas decreased SOD value in CP group increased in CP+RA group.

Our results show that CP causes histopathological damage in the rat kidney and that RA treatment relatively corrects these disorders caused by CP.

**Keywords:** *Rosmarinic acid, Cyclophosphamide, Toxicity, Kidney, Oxidative stress*

# How Does Nonsteroidal Anti-Inflammatory Ibuprofen Affect Neural Tube Development in Chick Embryos?

Dr. Esra Aslan<sup>1</sup>, Hilal Güzel<sup>2</sup>, Ph.D.

*Afyonkarahisar Health Sciences University, Faculty of Medicine, Department of Histology and Embryology, Afyonkarahisar, Turkey*<sup>1</sup>

*Afyonkarahisar Health Sciences University, Faculty of Medicine, Department of Anatomy, Afyonkarahisar, Turkey*<sup>2</sup>  
*dr\_esragul@hotmail.com*<sup>1</sup>; *hialgzl@hotmail.com*<sup>2</sup>

**Introduction:** Neural tube defects are heterogeneous and complex congenital anomalies of the central nervous system. The most common known causes are genetic factors and teratogenic drugs. All factors such as drugs, chemicals, infection or radiation that cause structural or functional disorders in the fetus when taken by the mother before or during pregnancy can have a "teratogenic" effect. Experimental early chick embryo model has a very traditional usage area in embryology. Since the neuronal and spinal developmental stages of the chick embryo are very close to the developmental stages of human embryos, it is a preferred subject group in studies on neural tube. Ibuprofen is one of the most commonly used non-steroidal anti-inflammatory drugs during pregnancy. Although the pregnancy category is in the C category, it has been found in the literature that developmental anomalies are observed. This study aimed to investigate the effects different doses of Ibuprofen on the neural tube development at 48 hours chick embryo models.

**Material and Methods:** 75 fertile, specific pathogen-free eggs were incubated for 28 hours and were divided into four groups of 15 eggs each (Group A: Control, Group B: 10mg/kg ibuprofen, Group C: 20mg/kg Ibuprofen, Group D: 40mg/kg Ibuprofen, and Group E: 80mg/kg Ibuprofen). Ibuprofen was administered via the sub-blastodermic route at this stage. Then the eggs were hand-turned 180° and placed in the incubator. Incubation was continued till the end of the 48<sup>th</sup> hour. All of the eggs were opened to 48<sup>th</sup> hour. All eggs were then opened and embryos were dissected from embryonic membranes. The samples were then transferred to petri dishes containing 10% formaldehyde solutions for histopathological study. Embryos stained with hematoxylin–eosin and the sections were examined using a light microscope.

**Results:** The effects of Ibuprofen on the embryo were correlated with the dose of Ibuprofen. It was determined that the use of increasing doses of Ibuprofen led to delayed of midline closure in early chick embryos. Somite numbers and crown-rump length were decreased in all experimental groups compared to the control group.

**Conclusion:** Neural tube defects are severe birth defects of the central nervous system that occur during embryonic development when the neural tube fails to close completely, leading to brain and spine anomalies that can lead to death or lifelong disability. This study showed that Ibuprofen delay neural tube development of chick embryo in a dose dependent manner. Results from animal studies may not reflect the circumstances in humans completely. The delayed neural tube mechanism of Ibuprofen isn't clear; therefore, studies of Ibuprofen should be supported with further researches.

**Keywords:** Ibuprofen, Chicken embryo, Neural tube